

Tachyon Condensation and Non-BPS D-branes in a Ramond-Ramond Plane Wave Background

Partha Mukhopadhyay

*Department of Applied Mathematics and Theoretical Physics
University of Cambridge
Wilberforce Road, Cambridge CB3 0WA, UK*

E-mail: P.Mukhopadhyay@damtp.cam.ac.uk

Abstract

We consider string theory in maximally supersymmetric type IIB plane wave background with constant five form Ramond-Ramond flux (RR plane wave). It is argued that there exists a universal sector of string configurations independent of the null coordinate x^- such that the space-time action evaluated at such a configuration is same in the RR plane wave and flat background. By naturally assuming its validity for the open strings we further argue that the D-branes extending along $x^\pm$ are universal in these two backgrounds. Moreover, a universal D-brane which is BPS in flat space must be tachyon-free in RR plane wave and a non-BPS D-brane should have a real tachyon whose potential is universal. Given the above observation we then proceed to describe open string theories for the non-BPS D-branes in RR plane wave. It is suggested that the light-cone Green-Schwarz fermions on the world-sheet satisfy certain bi-local boundary condition similar to that corresponding to flat space. We attempt a canonical quantisation with this boundary condition which gives rise to an open string spectrum similar to that in flat space - containing an R and NS sectors of states. In this process we encounter certain subtleties involving the R sector zero modes. Although the derivation of the R sector zero mode spectrum using the open string theory has not been completely settled, we suggest definite answer by requiring consistency with the relevant space-time interpretation. We finally generalise the above basic features of universality to all the exact pp-wave backgrounds with the same dilaton profile in a given string theory.

Contents

1	Introduction and summary	2
2	A universal sector of string configurations	6
3	Tachyon condensation and D-brane descent relation in RR plane wave	9
4	Non-BPS D-branes in RR Plane Wave	10
4.1	R Sector	12
4.2	NS Sector	17
4.3	Open string spectrum	17
5	Generalisation	19
A	Relevant features of even D-branes in RR plane wave	20
B	Open String Boundary Conditions in Green-Schwarz Formalism	21
B.1	Flat Background	21
B.2	RR Plane Wave Background	26
C	Identities	26
D	On the resolution of the zero-mode problem	27
E	A standard regularisation	33

1 Introduction and summary

Open string tachyon condensation¹ is an interesting and important subject of study in string theory. Its basic content is described by the well known conjectures due to Sen². Even though the associated physical picture is quite simple its complete string theoretic understanding is still lacking. The main obstacle is the mathematical complications of string theory which is also responsible for the fact that it has mainly been studied in flat space. It is therefore potentially interesting to study tachyon condensation in other geometric backgrounds in string theory³. In this regard one recalls that Sen argued [3]

¹By tachyon condensation we shall always refer to *open* string tachyon condensation in this paper.

²The reader is referred to the latest review [1] for the details of the developments in this subject and complete list of original references.

³See [2] for relevant studies in this direction.

the universality of tachyon potential in a certain class of superstring backgrounds which admit a symmetry under $(-1)^{F_L}$, where F_L is the number of space-time fermions that are left-moving on the world-sheet. In Sen's argument this symmetry was needed to define a coincident brane-anti-brane system. But there are other interesting backgrounds, namely the Ramond-Ramond (RR) backgrounds which do not admit such a symmetry. These backgrounds are important in string theory because of their role in the formulation of AdS/CFT correspondence [4] and the fact that string theory is poorly understood in such cases.

Among the non-trivial gravitational backgrounds plane waves are special as one gets more technical control on the quantisation problem in light-cone gauge. In this paper we focus on the particular example of type IIB maximally supersymmetric plane wave background with constant RR five form flux (RR plane wave) [5]. This is the simplest non-trivial example [6, 7, 8] that we would like to consider for our purpose. In light-cone Green-Schwarz (GS) formalism the world-sheet theory [6, 7] is simply a mass deformation of that corresponding to the flat background [9]. The mass parameter is given by a product of the strength of the background flux and the momentum p_- conjugate to the null coordinate x^- associated to the covariantly constant null Killing symmetry of the background. We point out that this indicates the following is true:

There exists a universal sector of string configurations independent of x^- such that the space-time action evaluated at such a configuration is same in RR plane wave and flat background. (1.1)

We shall present argument in favour of this conclusion in sec.2. Although our argument does not precisely specify what this universal sector is⁴, we argue that any scalar field configuration independent of x^- should be included in it. By naturally assuming the validity of the above statement for the open string theories we point out that it has the following consequence for the D-branes:

In type IIB string theory, for every D-brane in flat background extending along $x^\pm$ directions there exists a corresponding D-brane in RR plane wave background. (1.2)

This means that the D-branes extending along $x^\pm$ are *universal*. There are two categories of such D-branes, namely *even* and *odd*, depending on the dimensionality of the world-volume. As is well known, the even D-branes are half-BPS in flat space [10] while the

⁴This sector will be studied in more detail in [27]. See sec.5 for further comments on it.

odd ones break all the supersymmetries [11]. Although the detailed descriptions of such universal D-branes may be different in the two backgrounds, existence of the universal sector of string configurations predicts that the following must be true:

$$*A universal D-brane in RR plane wave is tachyon free if it is even and contains a real tachyon if it is odd. Moreover the tachyon potential is universal.* \quad (1.3)$$

D-branes in RR plane wave have already been analysed by many authors from various points of view [12, 13, 14, 15, 16]. According to the above classification these studies include only the even D-branes. Indeed all such D-branes have been found to be tachyon-free independently of their supersymmetries. One of the main goals of this paper is to develop the light-cone open string theories for the odd (non-BPS) D-branes. In flat space there are D-branes for which a space-like combination of the light-cone directions satisfies Dirichlet boundary condition while the orthogonal time-like combination satisfies Neumann boundary condition. Such D-branes can not be analysed in light-cone gauge and we do not discuss the analogues of such D-branes, if they exist, in RR plane wave.

Let us now discuss the non-BPS D-branes. In flat space the open string theories for such D-branes are not as straightforward as those corresponding to the even ones. In a manifestly supersymmetric formalism the complication lies in writing down the boundary condition for the space-time spinors. This problem was studied in light-cone gauge from both open and closed string points of view in [19, 20, 21]. In particular, it was found in [21] that the $SO(8)$ covariant open string boundary condition is actually bi-local which relates quadratic combinations of left and right moving GS fermions. Although the obvious question of how to deal with such a boundary condition was not completely settled, some progress was made. It was shown that this boundary condition leads to two sectors of open strings (see also [22]) namely, Ramond (R) and Neveu-Schwarz (NS) sectors. The resultant open string spectrum successfully passed through the world-sheet open-closed duality check⁵.

In this paper we show that all the valid open string boundary conditions for the kappa symmetry fixed type IIB GS superstrings in flat background are also valid in RR plane wave. This justifies us to use the similar bi-local boundary condition as discussed above

⁵Moreover, a proposal for how to compute disk amplitudes with all possible bulk insertions, but without any boundary insertion, was suggested [21] which produced correct results for the closed string one point functions. This boundary condition was generalised to its $SO(9, 1)$ covariant form and applied to pure spinor formalism [23] in [24, 25].

to define the non-BPS D-branes in RR plane wave. It is shown that, just like in flat space [21], there exist an R and NS sectors of open strings. However, there are certain additional subtleties involved in the present case. Usually a boundary condition enables us to solve for a set of degrees of freedom in terms of the rest. In absence of a linear boundary condition it is not *a priori* clear if the world-sheet lagrangian can be written down entirely in terms of the independent modes. We show that this can indeed be done almost for all the degrees of freedom except for the R sector zero modes - a complication which was absent in flat space.

We shall now briefly discuss this zero mode problem. Because of the mass term, the equations of motion dictate that the zero modes be either τ (world-sheet time) or σ (world-sheet space) dependent. It turns out that its only the τ dependent zero modes $S_0^{1a}(\tau)$ and $S_0^{2a}(\tau)$ for which one can possibly make sense of the proposed bi-local boundary condition. The problem is that while the two zero modes are related quadratically, the mass term in the lagrangian is linear in both and therefore can not be expressed in terms of any of the two. However, since this term does not contain any τ derivative, construction of the hamiltonian still goes through. Finally one arrives at a particular algebraic structure involving the constant zero modes S_0^a and $\tilde{S}_0^a$ which appear in the classical solutions for the τ dependent zero modes. This algebraic structure turns out to be quite unusual as S_0^a and $\tilde{S}_0^a$ are not linearly related. Although we are unable to settle the issue at this point, we argue that consistency with the relevant low-energy effective action requires us to have the same zero-mode spectrum as that in flat space.

In the NS sector the zero point energy does not cancel between the bosons and fermions because of broken supersymmetry. In flat space the standard zeta function regularisation gives the correct answer which produces a consistent spectrum for an interacting covariant field theory in space-time. The usual approach in the literature in the context of RR plane wave has been to consider the relevant Casimir energy which can be computed by using a generalised zeta function regularisation. But in this case we do not have an independent physical justification for this result. We remain inconclusive to this issue and leave its resolution as a future work.

The rest of the paper is organised as follows: We present our argument for the existence of the universal sector of string configurations in sec.2. The relevant consequences for the D-branes are discussed in sec.3. The open string theory for the GS fermions on a non-BPS D-brane in RR plane wave is discussed in sec.4. Finally we comment on certain

generalisations in sec.5. Several appendices contain necessary technical details.

2 A universal sector of string configurations

Here we shall argue in favour of the statement made in (1.1). The type IIB RR plane wave background is given by,

$$\begin{aligned} ds^2 &= G_{\mu\nu} dx^\mu dx^\nu , \\ &= 2dx^+ dx^- - f^2 x^I x^I dx^+ dx^+ + dx^I dx^I , \quad I = 1, \dots, 8 , \\ F_{+1234} &= F_{+5678} \propto f , \end{aligned} \tag{2.1}$$

where F is the RR five form field strength. Although f can be changed by properly scaling the light-cone coordinates the limit $f \rightarrow 0$ is special which corresponds to the flat background. String theory in this background has been solved in the standard light-cone gauge in [6, 7] which we refer the reader to for details. We shall follow the notations of [7] for our argument in this section and use them without any introduction.

The world-sheet theory in light-cone gauge is a mass deformation of that in flat background where the mass is given by⁶,

$$m = 2\pi\alpha' p_- f . \tag{2.2}$$

Although the bosonic sector of the theory maintains the complete transverse symmetry under $SO(8)$, the fermionic mass term breaks it to $SO(4) \times SO(4)$ as it is evident from the appearance of the matrix Π [7] which is the product of four gamma matrices along the directions: $1, \dots, 4$. This breaking of the global symmetry is expected due to the presence of the background flux.

Notice that the light-cone world-sheet theory is only sensitive to m which contains a product of p_- and f . This gives an indication that the $f \rightarrow 0$ limit is mimicked in the sector where the space-time fields are independent of the null coordinate x^- and therefore the background appears to be flat. To see this in a little more detail let us consider the space-time action in the background (2.1) written in terms of the Fourier transformed fields. A generic term in this action can be schematically written as,

$$T_{pw} = \int \delta^{10}(\sum_i p^i) \mathcal{A}(f, p_-^i, p_+^i, p_I^i) \prod_i \chi_i(p_-^i, p_+^i, p_I^i) , \tag{2.3}$$

⁶We use the notation p_- for the light-cone momentum which is same as p^+ used in [7].

where the integration is over all the momenta p^i . χ_i 's are the Fourier transformed fields. They carry space-time indices and can either be bosonic or fermionic. In general $\mathcal{A}$ is a function of all the momenta and the background parameter f . There are space-time index contractions among χ_i 's and $\mathcal{A}$ which are suppressed in this schematic expression. If $\mathcal{A}$ survives the $f \rightarrow 0$ limit then,

$$T_{flat} = \int \delta^{10}(\sum_i p^i) \mathcal{A}(0, p_-^i, p_+^i, p_I^i) \prod_i \chi_i(p_-^i, p_+^i, p_I^i) , \quad (2.4)$$

gives a term in the space-time action in flat background. If $\mathcal{A}$ does not survive the $f \rightarrow 0$ limit then it corresponds to certain interaction that is absent in flat space. If this limit has to be automatically mimicked in the sector where all the p_-^i 's are set to zero then we should have,

$$\mathcal{A}(f, p_-^i = 0, p_+^i, p_I^i) = \mathcal{A}(0, p_-^i = 0, p_+^i, p_I^i) . \quad (2.5)$$

Since we are assuming a smooth flat space limit for $\mathcal{A}$, it can be Taylor expanded and only positive powers of f appear. Also usually $\mathcal{A}$ follows some power law behaviour in terms of the momenta. With these assumptions the above equation implies,

$$\mathcal{A}(f, p_-^i, p_+^i, p_I^i) = \mathcal{A}'(m^i, p_-^i, p_+^i, p_I^i) , \quad (2.6)$$

where $\mathcal{A}'$ is another function and $m^i = 2\pi\alpha' p_-^i f$. This means that f always appears with a factor of p_-^i . $\mathcal{A}$ has to be computed using the world-sheet theory and we have already seen that this condition is indeed maintained in the light-cone gauge. Notice that the above argument does not precisely specify the universal sector that was referred to in (1.1). Although the required condition (2.5) or (2.6) is satisfied in light-cone gauge, it is not clear what should be the description of the universal sector outside the light-cone gauge. Work is in progress in [27] to answer such questions. Nevertheless a scalar field is certainly included in this sector and this is what we need for our argument on tachyon condensation in the next section.

We shall now discuss the symmetry algebra which is not same as that of the flat background. The kinematical and dynamical generators in the RR plane wave are [7]: P^+ , P^I , J^{+I} , J^{ij} ($i, j = 1, \dots, 4$), $J^{i'j'}$ ($i', j' = 5, \dots, 8$), Q^{+A} ($A = 1, 2$) and P^- , Q^{-A} respectively. In flat space the list includes certain extra kinematical generators, namely: J^{+-} , J^{-I} and $J^{ii'}$, which are missing in RR plane wave. Also the matching generators have different expressions in the two backgrounds. We therefore need to

1. account for the missing generators,
2. show that in expressions for the matching generators and algebra in RR plane wave f appears only as m .⁷

To account for the missing generators $J^{ii'}$ we notice that the matching generators J^{ij} and $J^{i'j'}$ do not depend on f (hence Π). Their construction can, in fact, be algebraically continued to incorporate $J^{ii'}$. The other missing generators J^{+-} and J^{-I} do not preserve the light-cone gauge and our analysis does not go beyond this gauge⁸. To analyse this we need a covariant setup with a precise knowledge of the universal sector in (1.1). But one can make the following comments for J^{+-} . This generates equal and opposite scaling of the coordinates x^+ and x^- . Certainly the background (2.1) is not invariant under such a transformation as it rescales the flux-strength f . But since $f \rightarrow 0$ limit is mimicked in the universal sector, such a transformation should appear as a symmetry in this sector.

To address the second point above we notice that the dynamical generators do depend on f and indeed the dependence is only through m . This, however, is not true of symmetry algebra - f explicitly appears in commutation relations. Schematically, f appears in the following way in such relations [7],

$$\begin{aligned} [P^-, P^I] &\sim f^2 J^{+I}, [P^-, Q^+] \sim f Q^+, [P^I, Q^-] \sim f Q^+, \\ \{Q^+, Q^-\} &\sim f J^{+I}, \{Q^-, Q^-\} \sim f J^{IJ}. \end{aligned} \quad (2.7)$$

But generators have additional p_- dependence:

$$\begin{aligned} P^+ &= p_-, P^- \sim \frac{1}{p_-}, J^{+I} \sim p_-, \\ Q^+ &\sim \sqrt{p_-}, Q^- \sim \frac{1}{\sqrt{p_-}}. \end{aligned} \quad (2.8)$$

These p_- dependence are such that they can be used in eqs.(2.7) to absorb all the f dependence and write the relevant commutation relations in terms of m .

⁷One should also expect that the Π dependence should go away in the limit $p_- \rightarrow 0$. This is automatically satisfied as $f \rightarrow 0$ is the flat space limit and therefore Π should always come with factors of f .

⁸I thank Peter van Nieuwenhuizen for asking pertinent questions on this.

3 Tachyon condensation and D-brane descent relation in RR plane wave

We shall now consider (1.1) in the context of D-branes and discuss its possible consequences for the tachyon condensation in RR plane wave. In particular, we shall argue below that the statement (1.2) is expected to be true.

To demonstrate the idea with the simplest example let us consider a non-BPS D-brane, Dp with p even, in type IIB string theory in flat background. Furthermore we orient it to extend along both the light-cone directions so that there are $(p-1)$ light-cone transverse directions (collectively denoted as $\vec{x}$) along the worldvolume. The worldvolume theory contains a real tachyon $T(x^+, x^-, \vec{x})$. This has a potential $V(T)$ so that the worldvolume action evaluated for a field configuration where the tachyon is set to a constant T and all the rest of the fields are set to zero is given by,

$$S_{flat}[T] = -\mathcal{V}^{(p+1)}V(T) , \quad (3.1)$$

where $\mathcal{V}^{(p+1)} = \int dx^+ dx^- d\vec{x}$ is the (infinite) volume of the worldvolume. $V(T)$ is an even function of T such that: (1) it has a maximum at $T = 0$ where the height of the potential is same as the tension of the D-brane, (2) it has non-perturbative minima at $|T| = T_0$ which correspond to the closed string vacuum as conjectured by Sen. It is natural to expect that S_{flat} can be obtained from a covariant action S_{cov} relevant to type IIB string theory by expanding it around the flat background. Given S_{cov} it should also be possible to expand it around the RR plane wave background (2.1) to obtain S_{pw} . As we have argued in the previous section that tachyon, being a scalar field, is included in the universal sector in (1.1). Therefore we must have,

$$S_{pw}[T] = -\mathcal{V}^{(p+1)}V(T) . \quad (3.2)$$

Now recall [1] that a suitable BPS $D(p-1)$ -brane in flat space can be viewed as a tachyonic kink solution $T(y)$ in the worldvolume theory described by S_{flat} .

$$T(y) = \begin{cases} \pm T_0 , & \text{as } y \rightarrow \pm\infty , \\ 0 , & \text{at } y = 0 , \end{cases} \quad (3.3)$$

where y is one of the light-cone transverse directions along the worldvolume of the parent D-brane that is transverse to the lower dimensional BPS D-brane. Notice that any solution

of the above kind is necessarily a solution of the theory described by the action S_{pw} as (1.1) applies to such configurations. Therefore one would expect this solution to correspond to an even D-brane in RR plane wave.

The basic argument used above also applies to the worldvolume theories on brane-anti-brane systems. Any D-brane in flat space that extends along x^- can be realised as a tachyonic solution that is independent of x^- in the worldvolume theory of sufficiently large number of brane-anti-brane systems or non-BPS D-branes (see [1] for the detailed discussion and references therein.). According to the above argument any such solution belongs to the universal sector of some worldvolume theory and therefore the corresponding D-brane is expected to exist in RR plane wave. This establishes a correspondence between the relevant D-branes in flat and the RR plane wave background as mentioned in (1.2). The only assumption in this argument is the existence of a covariant completion of any worldvolume theory that one obtains from open strings in flat space.

Although the above discussion implies that any universal D-brane is represented by the same worldvolume solution in flat background and RR plane wave, their detailed properties are in general different. For example, it is not guaranteed that a given universal D-brane preserves the same amount of supersymmetry in the two backgrounds. However, there are certain properties, such as those mentioned in (1.3), are guaranteed to be restricted by the existence of the universal sector. Referring the reader to (1.3), we comment that it is indeed true that all the even D-branes that have so far been discussed in the literature [12, 13, 14, 15, 16] are found to be tachyon-free. Given this discussion the next obvious question is: what is the open string theory for a non-BPS D-brane in RR plane wave that is predicted by universality? We suggest an answer to this question in the next section.

4 Non-BPS D-branes in RR Plane Wave

Here we shall discuss the open string theory for non-BPS D-branes in type IIB RR plane wave background in light-cone gauge. We have shown in appendix B that all the known open string boundary conditions in kappa gauge fixed GS action in flat background are also valid boundary conditions in RR plane wave⁹. This justifies considering the same

⁹Following [24] we have also written down the $SO(9, 1)$ covariant boundary condition for the space-time spinors for a non-BPS D-brane in flat space using the covariant GS action.

boundary conditions for the universal D-branes in both flat space and RR plane wave. Therefore in the bosonic sector we have the standard Neumann/Dirichlet boundary conditions - their analysis is quite standard and we shall not discuss them in detail. As in flat space the actual subtlety is involved in the space-time fermionic part [19, 20, 21] and therefore we shall focus our attention only to this sector in the following discussion.

The relevant world-sheet lagrangian is of the following form,

$$L = \beta \int_0^\pi d\sigma \left[(S^1 \partial_+ S^1) + (S^2 \partial_- S^2) - 2m (S^1 \Pi S^2) \right] , \quad (4.1)$$

where β is a constant and following the gamma matrix convention of [24] we have,

$$\Pi_{ab} = \sigma_{ab}^{1\cdots 4} = \sigma_{ab}^{5\cdots 8} . \quad (4.2)$$

The equations of motion read,

$$\partial_+ S^1 - m \Pi S^2 = 0 , \quad \partial_- S^2 + m \Pi S^1 = 0 , \quad (4.3)$$

The boundary condition, as discussed in appendix B, is given by [21],

$$S^{1a}(\tau, \sigma) S^{1b}(\tau', \sigma) = \mathcal{M}_{cd}^{ab} S^{2c}(\tau, \sigma) S^{2d}(\tau', \sigma) , \quad \text{at } \sigma = 0, \pi , \quad (4.4)$$

where the expression for $\mathcal{M}_{cd}^{ab}$ can be found in eq.(B.29). Since the fields are massive the equations of motion (4.3) mix S^1 and S^2 . Indeed using these equations of motion one can write down the following set of all possible boundary conditions,

$$S^{Aa}(\tau, \sigma) S^{Bb}(\tau', \sigma) = \mathcal{M}_{CDcd}^{ABab} S^{Cc}(\tau, \sigma) S^{Dd}(\tau', \sigma) , \quad \text{at } \sigma = 0, \pi , \quad (4.5)$$

where the non-zero entries for $\mathcal{M}_{CDcd}^{ABab}$ are given by:

$$\mathcal{M}_{22cd}^{11ab} = \mathcal{M}_{11cd}^{22ab} = \mathcal{M}_{cd}^{ab} , \quad \mathcal{M}_{21cd}^{12ab} = \mathcal{M}_{12cd}^{21ab} = - \left[\mathcal{M} \left(\frac{\Pi}{\Pi} \right) \right]_{cd}^{ab} , \quad (4.6)$$

where we have used the notation: $\left[\mathcal{M} \left(\frac{\Pi}{\Pi} \right) \right]_{cd}^{ab} = \mathcal{M}_{cc'}^{aa'} \Pi_{a'b} \Pi_{c'd}$. Since $\mathcal{M}_{cd}^{ab}$ has four indices Π_{ab} can be contracted in four different ways. The position of Π in the above notation indicates which index of $\mathcal{M}$ it is attached to¹⁰. The fields on the doubled surface (cylinder) can be defined similarly as in (B.21),

$$\mathcal{S}^{Aa}(\tau, \sigma) \mathcal{S}^{Bb}(\tau', \sigma') = \begin{cases} S^{Aa}(\tau, \sigma) S^{Bb}(\tau', \sigma') & 0 \leq \sigma, \sigma' \leq \pi , \\ \mathcal{M}_{CDcd}^{ABab} S^{Cc}(\tau, 2\pi - \sigma) S^{Dd}(\tau', 2\pi - \sigma') & \pi \leq \sigma, \sigma' \leq 2\pi . \end{cases} \quad (4.7)$$

¹⁰For example there is only one Π attached to $\mathcal{M}$ in $[\mathcal{M}(\Pi)]_{cd}^{ab}$ and three in $\left[\left(\frac{1}{\Pi} \right) \mathcal{M} \left(\frac{\Pi}{\Pi} \right) \right]_{cd}^{ab}$.

Using (4.7), (4.6) and (4.5) one can show,

$$\mathcal{S}^{Aa}(\tau, 2\pi)\mathcal{S}^{Bb}(\tau', 2\pi) = \mathcal{S}^{Aa}(\tau, 0)\mathcal{S}^{Bb}(\tau', 0) , \quad (4.8)$$

which implies,

$$\mathcal{S}^{Aa}(\tau, 2\pi) = \pm \mathcal{S}^{Aa}(\tau, 0) . \quad (4.9)$$

Therefore there exist two sectors (just like in flat space [21]): R and NS, corresponding to the above signs respectively. We shall now discuss the canonical quantisation for these two sectors separately below.

4.1 R Sector

R sector allows zero modes which can either be τ or σ dependent. As discussed in appendix A there are two classes of even D-branes, namely class I and class II, depending on whether the zero modes are τ or σ dependent respectively. The resulting algebraic structures are also different. We find it hard to use the present bi-local boundary condition to solve for dependent modes in terms of the independent ones in case of σ dependent zero modes. Below we shall show that with the τ dependent zero modes a consistent canonical quantisation for the open strings can be done with the proposed bi-local boundary condition leading to an algebraic structure which resembles more like the class I even D-branes.

We, therefore, start out with the following mode expansion:

$$\begin{aligned} S^1(\tau, \sigma) &= S_0^1(\tau) + \sum_{n \neq 0} c_n \left\{ S_n(\tau) e^{in\sigma} + i d_n \Pi \tilde{S}_n(\tau) e^{-in\sigma} \right\} , \\ S^2(\tau, \sigma) &= S_0^2(\tau) + \sum_{n \neq 0} c_n \left\{ \tilde{S}_n(\tau) e^{-in\sigma} - i d_n \Pi S_n(\tau) e^{in\sigma} \right\} , \end{aligned} \quad (4.10)$$

where,

$$c_n = \frac{1}{\sqrt{1 + d_n^2}} , \quad d_n = \frac{1}{m} (w_n - n) , \quad w_n = \text{sign}(n) \sqrt{n^2 + m^2} . \quad (4.11)$$

The classical equations of motion for the modes are given by,

$$\begin{aligned} \dot{S}_0^1(\tau) &= m \Pi S_0^2(\tau) , \quad \dot{S}_0^2(\tau) = -m \Pi S_0^1(\tau) , \\ \dot{S}_n(\tau) &= -i w_n S_n(\tau) , \quad \dot{\tilde{S}}_n(\tau) = -i w_n \tilde{S}_n(\tau) . \end{aligned} \quad (4.12)$$

All the degrees of freedom $S_0^1(\tau)$, $S_0^2(\tau)$, $S_n(\tau)$ and $\tilde{S}_n(\tau)$ are not independent because of the boundary condition (4.5). We shall now impose these conditions to determine the independent degrees of freedom. Naturally this task is more cumbersome than in the usual case of an even D-brane. Choosing $A = B = 1$ and $A = 1, B = 2$ in eq.(4.5) one arrives at two sets of relations involving these degrees of freedom. It turns out that there exists a set of independent solutions satisfying all such relations such that all these solutions relating various modes are independent of m . These solutions are bi-local and are given by,

$$\begin{aligned} S_0^{1a}(\tau)S_0^{1b}(\tau') &= \mathcal{M}_{cd}^{ab}S_0^{2c}(\tau)S_0^{2d}(\tau') , \\ S_0^{1a}(\tau)S_0^{2b}(\tau') &= -\left[\mathcal{M}\left(\frac{\Pi}{\Pi}\right)\right]_{cd}^{ab}S_0^{2c}(\tau)S_0^{1d}(\tau') . \end{aligned} \quad (4.13)$$

$$\begin{aligned} S_0^{1a}(\tau)S_n^b(\tau') &= \mathcal{M}_{cd}^{ab}S_0^{2c}(\tau)\tilde{S}_n^d(\tau') , \\ S_0^{1a}(\tau)\tilde{S}_n^b(\tau') &= -\left[\mathcal{M}\left(\frac{\Pi}{\Pi}\right)\right]_{cd}^{ab}S_0^{2c}(\tau)\tilde{S}_n^d(\tau') . \end{aligned} \quad (4.14)$$

$$\begin{aligned} S_n^a(\tau)S_{n'}^b(\tau') &= \mathcal{M}_{cd}^{ab}\tilde{S}_n^c(\tau)\tilde{S}_{n'}^d(\tau') , \\ S_n^a(\tau)\tilde{S}_{n'}^b(\tau') &= -\left[\mathcal{M}\left(\frac{\Pi}{\Pi}\right)\right]_{cd}^{ab}\tilde{S}_n^c(\tau)S_{n'}^d(\tau') . \end{aligned} \quad (4.15)$$

The above equations suggest that any expression that is quadratic in the set of modes $\{S^2(\tau), \tilde{S}_n(\tau)\}$ can be expressed entirely in terms of the set of modes $\{S_0^1(\tau), S_n(\tau)\}$ using the above conditions. We shall see that this will almost suffice for the present quantisation problem except for certain subtlety involving the zero modes which will be discussed as we go on.

We shall now attempt to compute the lagrangian (4.1) in terms of the independent modes using the relations (4.13, 4.14, 4.15). Substituting the expansion (4.10) in (4.1) one gets,

$$L = \beta(L_0 + L_1 + L_2) , \quad (4.16)$$

where, L_0 , L_1 and L_2 are the parts with the degree of non-zero modes being 0, 1 and 2 respectively. We find,

$$\frac{L_0}{2\pi} = \frac{1}{2} \left(S_0^1(\tau) \dot{S}_0^1(\tau) \right) + \frac{1}{2} \left(S_0^2(\tau) \dot{S}_0^2(\tau) \right) - m \left(S_0^1(\tau) \Pi S_0^2(\tau) \right) ,$$

$$= \left(S_0^1(\tau) \dot{S}_0^1(\tau) \right) - m \left(S_0^1(\tau) \Pi S_0^2(\tau) \right) , \quad (4.17)$$

where in the second step we have used the first equation in (4.13). Since this condition is bi-local in nature one can take τ -derivatives at the two points independently before taking the coincidence limit. Using the same technique with conditions (4.14) one finds,

$$L_1 = 0 . \quad (4.18)$$

Finally, using conditions (4.15) one obtains,

$$L_2 = 2\pi \sum_{n \neq 0} \left[\left(S_n(\tau) \dot{S}_{-n}(\tau) \right) - i w_n (S_n(\tau) S_{-n}(\tau)) \right] . \quad (4.19)$$

Notice that the lagrangian (4.16, 4.17, 4.18, 4.19) is almost entirely expressible in terms of the independent modes except for the mass term for the zero modes in (4.17) which is linear in $S_0^2(\tau)$. We can not express this term completely in terms of the independent modes as the conditions (4.13) are not factorisable to linear relations. Nevertheless we can smoothly proceed to construct the hamiltonian as this term does not contain any τ derivative. The canonically conjugate momenta are,

$$\Pi_0^a(\tau) = \frac{\partial L}{\partial \dot{S}_0^{1a}(\tau)} = 2\pi\beta S_0^{1a}(\tau) , \quad \Pi_n^a(\tau) = \frac{\partial L}{\partial \dot{S}_n^a(\tau)} = 2\pi\beta S_{-n}^a(\tau) . \quad (4.20)$$

The equal time canonical anti-commutators are found to be,

$$\{S_0^{1a}(\tau), S_0^{1b}(\tau)\} = \frac{i}{2\pi\beta} \delta^{ab} , \quad \{S_n^a(\tau), S_{n'}^b(\tau)\} = \frac{i}{2\pi\beta} \delta_{n+n',0} \delta^{ab} . \quad (4.21)$$

The hamiltonian turns out to be,

$$\begin{aligned} H &= H_0 - 4\pi i \beta \sum_{n>0} w_n (S_{-n} S_n) - 8 \sum_{n>0} w_n , \\ H_0 &= 2m\pi\beta \left(S_0^1(\tau) \Pi S_0^2(\tau) \right) , \end{aligned} \quad (4.22)$$

where the last term is the standard normal ordering constant and we have used the τ independent oscillators for the non-zero modes defined through the solutions of equations of motion:

$$S_n(\tau) = \exp(-i w_n \tau) S_n , \quad \forall n \neq 0 , \quad (4.23)$$

We shall now turn to the analysis of the zero modes below.

The zero-mode problem

The solutions to the zero mode equations of motion are given by,

$$S_0^1(\tau) = \cos(m\tau)S_0 + \sin(m\tau)\tilde{S}_0, \quad S_0^2(\tau) = \cos(m\tau)\Pi\tilde{S}_0 - \sin(m\tau)\Pi S_0. \quad (4.24)$$

Substituting these into eqs.(4.13) we find the following relations involving the τ -independent zero-modes:

$$\tilde{S}_0^a \tilde{S}_0^b = \widehat{\mathcal{M}}_{cd}^{ab} S_0^c S_0^d, \quad S_0^a \tilde{S}_0^b = -\widehat{\mathcal{M}}_{cd}^{ab} \tilde{S}_0^c S_0^d, \quad (4.25)$$

where $\widehat{\mathcal{M}}_{cd}^{ab}$ is obtained by replacing λ_I by $\hat{\lambda}_I$ in the expression of $\mathcal{M}_{cd}^{ab}$ in (B.29),

$$\widehat{\mathcal{M}}_{cd}^{ab} = \left[\left(\frac{1}{\Pi} \right) \mathcal{M} \left(\frac{1}{\Pi} \right) \right]_{cd}^{ab} = \mathcal{M}_{cd}^{ab}|_{\lambda_I \rightarrow \hat{\lambda}_I}, \quad (4.26)$$

where,

$$\hat{\lambda}_I = \lambda_I \beta_I, \quad (4.27)$$

and β_I is defined in eq.(C.5). The zero-mode equal-time anti-commutator in eqs.(4.21) implies,

$$\{S_0^a, S_0^b\} = \{\tilde{S}_0^a, \tilde{S}_0^b\} = \frac{i}{2\pi\beta} \delta^{ab}, \quad \{S_0^a, \tilde{S}_0^b\} = \Delta^{ab}, \quad (4.28)$$

such that,

$$\Delta^{ab} + \Delta^{ba} = 0. \quad (4.29)$$

The first set of equations in (4.28) are consistent with the first equation in (4.25). Using the first equation in (4.25) and (4.29) one finds the zero mode contribution to the hamiltonian to be,

$$H_0 = 2\pi m\beta(S_0\tilde{S}_0). \quad (4.30)$$

Given the algebraic structure in eqs.(4.25, 4.28) and (4.29) our final goal is to construct the zero-mode spectrum. This task is not straightforward as the relevant algebraic structure is quite unusual. Below we shall first try to understand the difficulty by comparing our case with that of the class I even D-branes and then find the expected result from an argument using the effective field theory on the worldvolume. Although it is not completely clear to us how to get this result from the open string theory, we discuss various arguments in appendix D for a possible resolution which reproduces this result.

Comparison with class I even D-branes

Certain relevant features of the even D-branes have been summarised in appendix A. Notice that the emergent algebraic structure is closer to that of the class I even D-branes than the ones belonging to class II. Equations (4.28, 4.29) and (4.30) also hold in case of class I D-branes, the only difference being that the relation between $\tilde{S}_0$ and S_0 is linear, unlike the ones in eqs.(4.25). This makes a big difference between the two cases. First of all, for class I D-branes H_0 can be expressed entirely in terms of the independent modes. Secondly, Δ^{ab} can be easily computed to be a c-number which enables one to construct the eigenstates of H_0 . Although the spectrum of H_0 is independent of Δ^{ab} , the H_0 eigenstates acquire certain angular momenta depending on Δ^{ab} [14]. On the contrary, in our case one can argue that Δ^{ab} can not be a non-zero c-number: Since Δ^{ab} is required to be anti-symmetric (see eq.(4.29)) it should be possible, if it is a c-number, to expand it in terms of the second rank $SO(8)$ gamma matrices. The manifest covariance of the whole construction would then require us to be able to single out two special directions. This can only be done for the class I even D-branes in this background because of their special $(n, n \pm 2)$ -kind of alignments (see appendix A). This is not true for the non-BPS D-branes and therefore there is no c-number choice for Δ^{ab} that is physically consistent in this case. This is effectively a variant of the statement that a linear relation does not exist between S_0^a and $\tilde{S}_0^a$. Moreover, notice that the second relation in (4.25), being quadratic, at most imposes certain constraints on Δ^{ab} without actually fixing it. Therefore the standard way of constructing the H_0 eigenstates does not go through.

What is expected?

The fact that $\tilde{S}_0^a$ could not be eliminated completely in our case and that Δ^{ab} does not get fixed might lead one to think that it is inconsistent to have these zero modes¹¹. This would then imply that there exists a unique ground state instead of the usual ones that transform as 8_V ($|I\rangle$) and 8_C ($|\dot{a}\rangle$) under $SO(8)$. But the existence of the universal sector of string configurations implies that these states should be present because of the following reason: Let us, for example, consider the vector states $|I\rangle$ - they correspond to the transverse components $A_I(x^+, x^-, \vec{x})$ of the $U(1)$ gauge field surviving in the light-cone gauge on the worldvolume. A configuration $A_I(x^+, \vec{x})$ independent of x^- is inside the universal

¹¹This was initially thought to be the case in [26].

sector and therefore should exist in RR plane wave. This is also true for the conjugate spinor states $|\dot{a}\rangle$. Given the above argument, the next question will be how H_0 acts on these ground states. The answer can be guessed in the following way: As mentioned in appendix A, for class I even D-branes H_0 breaks the degeneracy of the ground states which can be understood in the worldvolume theory as arising from a particular Chern-Simons coupling or its dimensional reduction. Such couplings are not present on the worldvolume of a non-BPS D-brane and therefore one would expect the degeneracy to be preserved in the present case. Therefore our conclusion is that the zero-mode spectrum is given by,

$$H_0|I\rangle = 0 \ , \quad H_0|\dot{a}\rangle = 0 \ . \quad (4.31)$$

4.2 NS Sector

The analysis in the NS sector is more straightforward as there are no zero modes. The mode expansions are,

$$\begin{aligned} S^1(\tau, \sigma) &= \sum_r c_r \left\{ S_r(\tau) e^{ir\sigma} + i d_r \Pi \tilde{S}_r(\tau) e^{-ir\tau} \right\} \ , \\ S^2(\tau, \sigma) &= \sum_r c_r \left\{ \tilde{S}_r(\tau) e^{-ir\sigma} - i d_r \Pi S_r(\tau) e^{ir\tau} \right\} \ , \end{aligned} \quad (4.32)$$

where r is an half-odd-integer and c_r , d_r and w_r are given by the same expressions as in eqs.(4.11) with n replaced by r . Classical equations of motion and the quadratic relations among the modes follow the equations for the non-zero modes given in eqs.(4.12) and (4.15) respectively with the similar replacement. Performing the similar computation as in the R sector one finally arrives at the hamiltonian:

$$H = -4\pi i \beta \sum_{r>0} w_r (S_{-r} S_r) - 8 \sum_{r>0} w_r \ , \quad (4.33)$$

in terms of the τ independent modes, defined similarly as in (4.23), which satisfy the following anti-commutation relation:

$$\{S_r^a, S_s^b\} = \frac{i}{2\pi\beta} \delta_{r+s,0} \delta^{ab} \ . \quad (4.34)$$

4.3 Open string spectrum

Based on the discussion in the previous subsections we are now in a position to spell out what the open string spectrum is. For a non-BPS D-brane sitting at the origin of the

transverse directions the light-cone hamiltonian in R and NS sectors are given by,

$$H^R = H_B + H_F^R, \quad H^{NS} = H_B + H_F^{NS}, \quad (4.35)$$

respectively where the bosonic part is given by,

$$2\alpha' p_- H_B = \frac{m}{\pi} a^{I\parallel\dagger} a^{I\parallel} + \frac{m}{2\pi} (p-1) + \sum_{n>0} \alpha_{-n}^I \alpha_n^I. \quad (4.36)$$

Recall the expression for m given in eq.(2.2). $I_{\parallel}$ runs over the light-cone transverse directions that are parallel to the world-volume. The oscillators $a^{I\parallel}$ are constructed out of the bosonic zero modes and α_n^I 's are the standard bosonic non-zero modes. The commutation relations are:

$$[a^{I\parallel}, a^{J\parallel\dagger}] = \delta^{IJ} J_{\parallel}, \quad [\alpha_n^I, \alpha_{n'}^J] = \delta^{IJ} w_n \delta_{n+n',0}. \quad (4.37)$$

The second term on the right hand side of eq.(4.36) is simply the normal ordering constant coming from the bosonic zero mode oscillators. The ground states in the R sector are $|I\rangle$ and $|\dot{a}\rangle$ which are annihilated by $a^{I\parallel}$, α_n^I and S_n^a for $n > 0$. The unique ground state $|0\rangle$ in the NS sector is, as usual, defined to be annihilated by the same bosonic oscillators and S_r^a for $r > 0$. As argued in sec.(4.1) the fermionic zero modes do not contribute to the hamiltonian in the R sector. The non-zero mode parts in the R and NS sectors are given by,

$$2\alpha' p_- H_F^R = \sum_{n>0} w_n S_{-n}^a S_n^a, \quad 2\alpha' p_- H_F^{NS} = \sum_{r>0} w_r S_{-r}^a S_r^a + h_0(m), \quad (4.38)$$

where,

$$w_k = \sqrt{k^2 + \left(\frac{m}{2\pi}\right)^2}, \quad (4.39)$$

with k being either integer or half-odd-integer. The fermionic oscillators satisfy the relevant anti-commutators of eqs.(4.21, 4.34) with $\beta = i/2\pi$. The zero point energy for the non-zero modes cancel between bosons and fermions in the R sector. However, this does not happen in the NS sector where it is given by,

$$h_0(m) = 4 \left(\sum_{n>0} w_n - \sum_{r>0} w_r \right). \quad (4.40)$$

This contains divergent sums and usually one regularises this following standard methods. This has been done in appendix E and the result is a non-trivial function of m involving certain modified Bessel functions. But as mentioned before, unlike in the case of flat space we do not yet have an independent physical understanding of this result. It would be interesting to explore this issue in future.

5 Generalisation

Plane wave backgrounds are special cases of a more general backgrounds called pp-waves which admit a covariantly constant null Killing symmetry. In all such backgrounds the standard light-cone gauge can be fixed [30]. Quite generically the world-sheet theories in light-cone gauge involve a mass scale set by p_- [31] such that the light-cone Hamiltonian takes the following form:

$$p_- H_{pp} = p_- H_{flat} + \mathcal{I}(p_-) , \quad (5.1)$$

where $\mathcal{I}(p_-)$ is a collection of interaction terms representing the deformation from flat space. This interaction term depends on p_- in such a way that,

$$\lim_{p_- \rightarrow 0} \mathcal{I}(p_-) \rightarrow 0 . \quad (5.2)$$

This is clearly the case for the RR plane wave background, considered in detail in this paper. This is also true for some of the cases explicitly studied in the literature [32, 33]. This raises the following question:

Does the universality hold in all the pp-wave backgrounds?

This question will be addressed in [27]. The answer turns out to be yes (which implies that the space-time action will be evaluated to be same around the pp-wave backgrounds) only when all the scalars (for example dilaton) that might be switched on in the background take the same profile¹². Notice that dilaton is constant in both the flat and RR plane wave backgrounds studied here. This can further be generalised to non-trivial transverse

¹²For the model studied in [33] the interaction is of special type such that eq.(5.2) does not hold in light-cone gauge. But the universality might still work as the interaction term does drop off in the linearised equation of motion in space-time for configurations which do not depend on x^- (see eq.(5.19) in [33]).

spaces or to cases where the transverse space is consistently replaced by a conformal field theory.

Acknowledgement

I am thankful to Sumit R. Das, Matthias R. Gaberdiel, Michael B. Green, Martin Kruczenski, Juan Maldacena, Ignacio Navarro, Alfred D. Shapere and Aninda Sinha for useful discussion. A preliminary version of this work was presented at EFI, University of Chicago; YITP, Stony Brook, Kentucky University, Great Lakes Strings Conference 2006, MCTP, University of Michigan in March 2006 and Purdue University in October 2006. I thank the High Energy Physics Theory groups at the above mentioned places and the organisers of the Michigan conference for their interest. I am also grateful to Michael B. Green for reading an earlier version of the manuscript and a following very useful discussion and Sumit R. Das and Alfred D. Shapere for their continuous encouragement and for supporting visits to Kentucky University where parts of this work were done. This work was financially supported by PPARC.

A Relevant features of even D-branes in RR plane wave

Here we shall briefly review certain features of the open strings on even D-branes in RR plane wave background that are relevant to the R-sector zero mode problem for the non-BPS D-branes discussed in this paper. Because of the mass term the zero modes can either be τ or σ dependent. This leads to two kinds of zero mode structures for the even D-branes. The τ dependent zero modes end up giving a non-zero contribution to the light-cone hamiltonian which breaks the degeneracy of the 8_v and 8_c ground states. These D-branes have been called Class I [13] or D_- [14] branes in the literature. This lifting of the degeneracy can be understood in the following way. All these D-branes are either D7 or their dimensional reductions. The orientation of such a D-brane is usually denoted as $(n, n \pm 2)$, which implies that in addition to $x^\pm$ the brane is extended along n directions out of $x^1, \dots, x^4$ and $(n \pm 2)$ directions out of $x^5, \dots, x^8$. Notice that the maximum worldvolume dimensionality of such a D-brane is 8 corresponding to a D7-brane. A D7-brane is special in this background in the following sense. Its eight dimensional worldvolume contains a Chern-Simons (CS) coupling with the background RR 4-form potential which is quadratic in the worldvolume field strength [12]. Hence this introduces

certain extra term in the linearised equations of motion for the transverse components of the worldvolume vector potential which is responsible for the lifting of degeneracy of the open string ground states in 8_V . The supersymmetric version of this explains why the degeneracy of the 8_C states should also be broken. None of the other even D-branes (called Class II or D_+ branes) contains such a CS coupling. The open string zero modes for such a D-brane is σ dependent which do not contribute to the light-cone hamiltonian preserving the degeneracy of the 16 ground states.

B Open String Boundary Conditions in Green-Schwarz Formalism

Here we shall show that the open string boundary conditions for the BPS and non-BPS D-branes of type IIB string theory in flat background that can be analysed using kappa gauge fixed GS action are also valid boundary conditions in the RR plane-wave background. Below we shall first review the flat background where our discussion will be both in the pre and post kappa gauge fixed theory. Later we shall show why the kappa gauge fixed boundary conditions are also valid in RR plane wave.

B.1 Flat Background

After gauge fixing the world-sheet metric: $g_{ab} = \eta_{ab}$ the GS world-sheet action takes the following form:

$$\begin{aligned} S &\propto S_1 + S_2 , \\ S_1 &= -\frac{1}{2} \int d^2\sigma \eta^{ab} \eta_{\mu\nu} \left(\partial_a X^\mu - i(\theta^A \bar{\gamma}^\mu \partial_a \theta^A) \right) \left(\partial_b X^\nu - i(\theta^B \bar{\gamma}^\nu \partial_b \theta^B) \right) , \\ S_2 &= \int d^2\sigma \epsilon^{ab} \eta_{\mu\nu} \left[-i \partial_a X^\mu \left\{ (\theta^1 \bar{\gamma}^\nu \partial_b \theta^1) - (\theta^2 \bar{\gamma}^\nu \partial_b \theta^2) \right\} + (\theta^1 \bar{\gamma}^\mu \partial_a \theta^1) (\theta^2 \bar{\gamma}^\nu \partial_b \theta^2) \right] , \end{aligned} \quad (B.1)$$

where $a, b = 0, 1$, $\sigma^0 = \tau$, $\sigma^1 = \sigma$, ϵ^{ab} is the two dimensional Levi-Civita symbol with $\epsilon^{01} = 1$. $A = 1, 2$ and we follow the convention of [24] for the 10-dimensional spinor and the gamma matrices. The boundary terms obtained upon varying the basic fields are given by,

$$\delta S_{boundary} \propto \int d\tau [-T_1 + iT_2 - T_3 + T_4 + iT_5]_0^\pi , \quad (B.2)$$

where $[A]_0^\pi = A(\tau, \sigma = \pi) - A(\tau, \sigma = 0)$ and

$$\begin{aligned}
T_1 &= \delta X^\mu \partial_\sigma X_\mu , \\
T_2 &= \partial_\sigma X^\mu \left\{ (\theta^1 \bar{\gamma}_\mu \delta \theta^1) + (\theta^2 \bar{\gamma}_\mu \delta \theta^2) \right\} - \partial_\tau X^\mu \left\{ (\theta^1 \bar{\gamma}_\mu \delta \theta^1) - (\theta^2 \bar{\gamma}_\mu \delta \theta^2) \right\} , \\
T_3 &= (\theta^1 \bar{\gamma}_\mu \delta \theta^1) (\theta^2 \bar{\gamma}^\mu \partial_\tau \theta^2) - (\theta^2 \bar{\gamma}^\mu \delta \theta^2) (\theta^1 \bar{\gamma}_\mu \partial_\tau \theta^1) , \\
T_4 &= \left\{ (\theta^1 \bar{\gamma}_\mu \delta \theta^1) + (\theta^2 \bar{\gamma}_\mu \delta \theta^2) \right\} \left\{ (\theta^1 \bar{\gamma}^\mu \partial_\sigma \theta^1) + (\theta^2 \bar{\gamma}^\mu \partial_\sigma \theta^2) \right\} , \\
T_5 &= \delta X^\mu \left\{ (\theta^1 \bar{\gamma}_\mu \partial_\sigma \theta^1) + (\theta^2 \bar{\gamma}_\mu \partial_\sigma \theta^2) \right\} + \delta X^\mu \left\{ (\theta^1 \bar{\gamma}_\mu \partial_\tau \theta^1) - (\theta^2 \bar{\gamma}_\mu \partial_\tau \theta^2) \right\} . \quad (B.3)
\end{aligned}$$

Components of the stress-tensor are given by,

$$\begin{aligned}
T_{\tau\tau} &= T_{\sigma\sigma} = \frac{1}{2} (\partial_\tau X^\mu \partial_\tau X_\mu + \partial_\sigma X^\mu \partial_\sigma X_\mu) + i \partial_\tau X_\mu \sum_A \left(\theta^A \bar{\gamma}^\mu \partial_\tau \theta^A \right) + i \partial_\sigma \sum_A \left(\theta^A \bar{\gamma}^\mu \partial_\sigma \theta^A \right) , \\
T_{\tau\sigma} &= \partial_\tau X^\mu \partial_\sigma X_\mu + i \partial_\tau X_\mu \sum_A \left(\theta^A \bar{\gamma}^\mu \partial_\sigma \theta^A \right) + i \partial_\sigma X_\mu \sum_A \left(\theta^A \bar{\gamma}^\mu \partial_\tau \theta^A \right) . \quad (B.4)
\end{aligned}$$

Although the Virasoro constraint $T_{\tau\tau} = 0$ imposes restriction on the physical sates, in order for the energy eigenvalue to be independent of τ we need to satisfy identically,

$$T_{\tau\sigma} = 0 , \quad \text{at } \sigma = 0, \pi . \quad (B.5)$$

We shall now discuss open string boundary conditions corresponding to BPS and non-BPS D-branes for which the variation in (B.2) vanishes and eq.(B.5) is satisfied identically. The standard Neumann and Dirichlet boundary conditions for the space-time vectors are given by,

$$\begin{aligned}
\text{Neumann:} \quad & \partial_\sigma X^\mu(\tau, \sigma) = 0 , \quad \text{at } \sigma = 0, \pi , \\
\text{Dirichlet:} \quad & \delta X^\mu(\tau, \sigma) = 0 \text{ or } \partial_\tau X^\mu(\tau, \sigma) = 0 , \text{ at } \sigma = 0, \pi . \quad (B.6)
\end{aligned}$$

Due to this T_1 vanishes in equation (B.2). Boundary conditions for the space-time spinors are different for BPS and non-BPS D-branes which will be discussed separately below:

BPS D-Branes

For a BPS D-brane the relevant boundary condition is given by [17, 18],

$$\begin{aligned}
\theta^1(\tau, \sigma) &= M^S \theta^2(\tau, \sigma) , \quad \text{at } \sigma = 0, \pi , \\
M^S &= \gamma^{\mu_1 \dots \mu_{p+1}} , \quad (B.7)
\end{aligned}$$

where $\mu_1, \dots, \mu_{p+1}$ correspond to the world-volume directions of a Dp -brane (p odd). Boundary conditions involving derivatives read,

$$\partial_\tau \theta^1(\tau, \sigma) = M^S \partial_\tau \theta^2(\tau, \sigma) , \quad \partial_\sigma \theta^1(\tau, \sigma) = -M^S \partial_\sigma \theta^2(\tau, \sigma) , \quad \text{at } \sigma = 0, \pi . \quad (\text{B.8})$$

One way to see the origin of the sign in the second equation is the following. Define a spinor $\Theta(\tau, \sigma)$ on the cylinder: $-\infty \leq \tau \leq \infty$, $0 \leq \sigma \leq 2\pi$ by the standard doubling trick:

$$\Theta(\tau, \sigma) = \begin{cases} \theta^1(\tau, \sigma) , & 0 \leq \sigma \leq \pi , \\ M^S \theta^2(\tau, 2\pi - \sigma) , & \pi \leq \sigma \leq 2\pi . \end{cases} \quad (\text{B.9})$$

Then for small ϵ ,

$$\partial_\sigma \Theta(\tau, \sigma)|_{\sigma=\pi-\epsilon} = \partial_\sigma \theta^1(\tau, \sigma)|_{\sigma=\pi-\epsilon} . \quad (\text{B.10})$$

On the other hand,

$$\begin{aligned} \partial_\sigma \Theta(\tau, \sigma)|_{\sigma=\pi+\epsilon} &= M^S \partial_\sigma \theta^2(\tau, 2\pi - \sigma)|_{\sigma=\pi+\epsilon} , \\ &= -M^S \partial_{\tilde{\sigma}} \theta^2(\tau, \tilde{\sigma})|_{\tilde{\sigma}=\pi-\epsilon} . \end{aligned} \quad (\text{B.11})$$

Taking the limit $\epsilon \rightarrow 0$ on the right hand sides of eqs.(B.10), (B.11) and equating them one gets the second equation in (B.8). Using the boundary conditions (B.7, B.8) and the relations,

$$(M^S)^T \bar{\gamma}^\mu M^S = - (M^V)^\mu{}_\nu \bar{\gamma}^\nu , \quad (\text{B.12})$$

$$\left. \begin{aligned} &\left[\delta^\mu{}_\nu - (M^V)^\mu{}_\nu \right] \partial_\sigma X^\nu = 0 , \\ &\left[\delta^\mu{}_\nu + (M^V)^\mu{}_\nu \right] \partial_\tau X^\nu = 0 , \text{ or } \left[\delta^\mu{}_\nu + (M^V)^\mu{}_\nu \right] \delta X^\nu = 0 , \end{aligned} \right\} \text{at } \sigma = 0, \pi , \quad (\text{B.13})$$

where,

$$(M^V)^\mu{}_\nu = \begin{cases} -\delta^\mu{}_\nu , & \text{for } X^\mu \text{ Neumann} , \\ \delta^\mu{}_\nu , & \text{for } X^\mu \text{ Dirichlet} . \end{cases} \quad (\text{B.14})$$

one can show that all the rest of the terms in the variation (B.2) vanish. Similarly using the boundary conditions (B.6), (B.7) and (B.8) and the relation (B.13) one can show that eq.(B.5) is satisfied identically.

Non-BPS D-Branes

For non-BPS D-branes we propose, following [24], the following bi-local boundary condition for the space-time spinors:

$$\theta^{1\alpha}(\tau, \sigma) \theta^{1\beta}(\tau', \sigma) = \mathcal{M}_{\gamma\delta}^{\alpha\beta} \theta^{2\gamma}(\tau, \sigma) \theta^{2\delta}(\tau', \sigma) , \quad \text{at } \sigma = 0, \pi , \quad (\text{B.15})$$

where,

$$\begin{aligned} \mathcal{M}_{\gamma\delta}^{\alpha\beta} = & - \left[\frac{1}{16} \gamma_{\mu}^{\alpha\beta} (M^V)^{\mu}_{\nu} \bar{\gamma}_{\gamma\delta}^{\nu} + \frac{1}{16 \times 3!} \gamma_{\mu_1 \dots \mu_3}^{\alpha\beta} (M^V)^{\mu_1}_{\nu_1} \dots (M^V)^{\mu_3}_{\nu_3} \bar{\gamma}_{\gamma\delta}^{\nu_1 \dots \nu_3} \right. \\ & \left. + \frac{1}{16 \times 5!} \sum_{\mu_1, \dots, \mu_5 \in \mathcal{K}^{(5)}} \gamma_{\mu_1 \dots \mu_5}^{\alpha\beta} (M^V)^{\mu_1}_{\nu_1} \dots (M^V)^{\mu_5}_{\nu_5} \bar{\gamma}_{\gamma\delta}^{\nu_1 \dots \nu_5} \right] . \end{aligned} \quad (\text{B.16})$$

The definition of the set $\mathcal{K}^{(5)}$ can be found in [25] (see discussion below eq.(2.3) in [25]). We shall now argue that all the terms $[T_2]_0^{\pi}, \dots, [T_5]_0^{\pi}$ on the right hand side of eq.(B.2) vanish. Considering the term T_2 one finds,

$$[T_2]_0^{\pi} = \left[\left\{ \partial_{\sigma} X_{\mu} \left(\delta^{\mu}_{\nu} - (M^V)^{\mu}_{\nu} \right) - \partial_{\tau} X_{\mu} \left(\delta^{\mu}_{\nu} + (M^V)^{\mu}_{\nu} \right) \right\} (\theta^2 \bar{\gamma}^{\nu} \delta \theta^2) \right]_0^{\pi} = 0 , \quad (\text{B.17})$$

where we have used

$$\left(\theta^1(\tau, \sigma) \bar{\gamma}^{\mu} \theta^1(\tau', \sigma) \right) + M^{\mu}_{\nu} \left(\theta^2(\tau, \sigma) \bar{\gamma}^{\nu} \theta^2(\tau', \sigma) \right) = 0 , \quad \text{at } \sigma = 0, \pi , \quad (\text{B.18})$$

which can be obtained from the boundary condition (B.15). To treat the rest of the terms we need to know the boundary conditions which involve variation and derivatives of the space-time spinors. Since the boundary condition (B.18) is bi-local we can vary the fields at the two points independently. This gives,

$$\left(\theta^1(\tau, \sigma) \bar{\gamma}^{\mu} \delta \theta^1(\tau', \sigma) \right) + M^{\mu}_{\nu} \left(\theta^2(\tau, \sigma) \bar{\gamma}^{\nu} \delta \theta^2(\tau', \sigma) \right) = 0 , \quad \text{at } \sigma = 0, \pi . \quad (\text{B.19})$$

One can also apply τ -derivatives at the two points independently. Therefore,

$$\left(\theta^1(\tau, \sigma) \bar{\gamma}^{\mu} \partial_{\tau} \theta^1(\tau', \sigma) \right) + M^{\mu}_{\nu} \left(\theta^2(\tau, \sigma) \bar{\gamma}^{\nu} \partial_{\tau} \theta^2(\tau', \sigma) \right) = 0 , \quad \text{at } \sigma = 0, \pi . \quad (\text{B.20})$$

Using these it is straightforward to show that $[T_3]_0^{\pi} = 0$. To obtain the boundary condition involving σ -derivative we first adopt a suitable doubling trick following [21]:

$$\Theta^{1\alpha}(\tau, \sigma) \Theta^{1\beta}(\tau', \sigma') = \begin{cases} \theta^{1\alpha}(\tau, \sigma) \theta^{1\beta}(\tau', \sigma') & 0 \leq \sigma, \sigma' \leq \pi , \\ \mathcal{M}_{\gamma\delta}^{\alpha\beta} \theta^{2\gamma}(\tau, 2\pi - \sigma) \theta^{2\delta}(\tau', 2\pi - \sigma') & \pi \leq \sigma, \sigma' \leq 2\pi . \end{cases} \quad (\text{B.21})$$

Then using the similar reasoning as done in the BPS-case one can show:

$$\left(\theta^1(\tau, \sigma) \bar{\gamma}^{\mu} \partial_{\sigma} \theta^1(\tau', \sigma) \right) - M^{\mu}_{\nu} \left(\theta^2(\tau, \sigma) \bar{\gamma}^{\nu} \partial_{\sigma} \theta^2(\tau', \sigma) \right) = 0 , \quad \text{at } \sigma = 0, \pi . \quad (\text{B.22})$$

Using this, the boundary conditions (B.20) and (B.13) one can show $[T_4]_0^\pi = 0$ and $[T_5]_0^\pi = 0$. This shows that for the boundary conditions (B.6) and (B.15) the boundary terms (B.2) in the variation of the world-sheet action are zero. Also using eqs.(B.6), (B.20), (B.22) and (B.13) one can establish that eq.(B.5) is satisfied identically.

D-Branes in Kappa Gauge Fixed Theory

If we fix the fermionic kappa symmetry by requiring:

$$\bar{\gamma}^+ \theta^A = 0 \ , \Rightarrow \theta^A = c \begin{pmatrix} S^a \\ 0 \end{pmatrix} \ , \quad (\text{B.23})$$

then we have,

$$\begin{aligned} T_2 &= \sqrt{2}c^2 \left[\partial_\sigma X^+ \left\{ (S^1 \delta S^1) + (S^2 \delta S^2) \right\} - \partial_\tau X^+ \left\{ (S^1 \delta S^1) - (S^2 \delta S^2) \right\} \right] \ , \\ T_3 &= 0 \ , \\ T_4 &= 0 \ , \\ T_5 &= \sqrt{2}c^2 \left[\delta X^+ \left\{ (S^1 \partial_\sigma S^1) + (S^2 \partial_\sigma S^2) \right\} + \delta X^+ \left\{ (S^1 \partial_\tau S^1) - (S^2 \partial_\tau S^2) \right\} \right] \ , \end{aligned} \quad (\text{B.24})$$

and

$$\begin{aligned} T_{\tau\sigma} &= \partial_\tau X^+ \partial_\sigma X^- + \partial_\sigma X^+ \partial_\tau X^- + \partial_\tau X^I \partial_\sigma X^I \\ &\quad + \sqrt{2}ic^2 \partial_\tau X^+ \sum_A (S^A \partial_\sigma S^A) + \sqrt{2}ic^2 \partial_\sigma X^+ \sum_A (S^A \partial_\tau S^A) \ . \end{aligned} \quad (\text{B.25})$$

For all the D-branes that can be described in this gauge, both the light-cone coordinates $X^\pm$ should satisfy Neumann boundary condition¹³:

$$\partial_\sigma X^\pm(\tau, \sigma) = 0 \ , \quad \text{at } \sigma = 0, \pi \ . \quad (\text{B.26})$$

Boundary condition for the space-time spinors for a BPS D-brane reads:

$$\begin{aligned} S^{1a}(\tau, \sigma) &= \mathcal{M}_{ab}^S S^{2b}(\tau, \sigma) \ , \quad \text{at } \sigma = 0, \pi \ , \\ \mathcal{M}_{ab}^S &= \left(\sigma^{I_1 I_2 \dots I_{p-1}} \right)_{ab} \ , \end{aligned} \quad (\text{B.27})$$

where $X^{I_1}, \dots, X^{I_{p-1}}$ are the Neumann directions besides $X^\pm$. For a non-BPS D-brane one gets,

$$S^{1a}(\tau, \sigma) S^{1b}(\tau', \sigma) = \mathcal{M}_{cd}^{ab} S^{2c}(\tau, \sigma) S^{2d}(\tau', \sigma) \ , \quad (\text{B.28})$$

¹³In order for the first term in the expression for T_2 in eqs.(B.24) to vanish at the boundary we need X^+ to satisfy Neumann boundary condition. Then in order for the first term in eq.(B.25) to vanish at the boundary we need X^- also to satisfy Neumann boundary condition.

where,

$$\mathcal{M}_{cd}^{ab} = \frac{1}{8}\delta_{ab}\delta_{cd} + \frac{1}{16}\sum_{IJ}\lambda_I\lambda_J\sigma_{ab}^{IJ}\sigma_{cd}^{IJ} + \frac{1}{192}\sum_{IJKL\in\mathcal{K}^{(4)}}\lambda_I\lambda_J\lambda_K\lambda_L\sigma_{ab}^{IJKL}\sigma_{cd}^{IJKL} \quad (\text{B.29})$$

As explained in [21], the summation over the vector indices in the last term is restricted and a definition for the set $\mathcal{K}^{(4)}$ can be found, for example, in [25] (see discussion below eq.(4.9)). λ_I is the eigenvalue of M^V in eq.(B.14) along the light-cone transverse direction x^I .

B.2 RR Plane Wave Background

In case of the RR plane wave background the boundary terms in the variation of the world-sheet action read:

$$\delta S_{pw}^{boundary} = \delta S^{boundary} + \delta(\Delta S) , \quad (\text{B.30})$$

where $\delta S^{boundary}$ is given in eq.(B.2) with T_1 given in (B.3) and the $T_2, \dots, T_5$ given in (B.24) and,

$$\delta(\Delta S) \propto \int d\tau \left[\mu^2 \vec{X}^2 \delta X^+ \partial_\sigma X^+ + 4\sqrt{2}ic^2 \mu \delta X^+ \partial_\sigma X^+ (S^1 \Pi S^2) \right]_0^\pi . \quad (\text{B.31})$$

The off-diagonal component of the stress-tensor is given by,

$$(T_{pw})_{\tau\sigma} = T_{\tau\sigma} + \Delta T_{\tau\sigma} , \quad (\text{B.32})$$

where $T_{\tau\sigma}$ is given in eq.(B.25) and,

$$\Delta T_{\tau\sigma} = -\mu^2 \vec{X}^2 \partial_\tau X^+ \partial_\sigma X^+ - 4\sqrt{2}ic^2 \mu \partial_\tau X^+ \partial_\sigma X^+ (S^1 \Pi S^2) . \quad (\text{B.33})$$

Notice that all the terms in $\delta(\Delta S)$ and $\Delta T_{\tau\sigma}$ contain $\partial_\sigma X^+$. Since X^+ satisfies Neumann boundary condition both $\delta(\Delta S)$ and $\Delta T_{\tau\sigma}$ at the boundary are zero. This shows that all the open string boundary conditions in flat space are also valid boundary conditions in the RR plane wave.

C Identities

Here we present some useful identities that are required for various computations done in this paper. Using the following completeness relation,

$$\frac{1}{8}\delta_{ab}\delta_{ef} + \frac{1}{16}\sum_{I,J}\sigma_{ab}^{IJ}\sigma_{ef}^{IJ} + \frac{1}{384}\sum_{I,J,K,L}\sigma_{ab}^{IJKL}\sigma_{ef}^{IJKL} = \delta_{ae}\delta_{bf} , \quad (\text{C.1})$$

one can show,

$$\mathcal{M}_{cd}^{ab} \mathcal{M}_{ef}^{cd} = \delta_{ae} \delta_{bf} . \quad (\text{C.2})$$

We also have,

$$\mathcal{M}_{cd}^{ab} = \mathcal{M}_{dc}^{ba} , \quad (\text{C.3})$$

$$\Pi^T = \Pi , \quad \Pi \sigma^{IJ\dots} = (\beta_I \beta_J \dots) \sigma^{IJ\dots} \Pi , \quad (\text{C.4})$$

where in the last equation of (C.4) product of an even number of sigma matrices has been considered and,

$$\beta_I = \begin{cases} 1 & \text{if } I \in \{1, 2, 3, 4\} \\ -1 & \text{otherwise} . \end{cases} \quad (\text{C.5})$$

Using the above relations one can prove the following identities:

$$\left[\left(\frac{\Pi^{\epsilon_1}}{\Pi^{\epsilon_2}} \right) \mathcal{M} \left(\frac{\Pi^{\epsilon_3}}{\Pi^{\epsilon_4}} \right) \right]_{cd}^{ab} = \left[\left(\frac{\Pi^{1-\epsilon_1}}{\Pi^{1-\epsilon_2}} \right) \mathcal{M} \left(\frac{\Pi^{1-\epsilon_3}}{\Pi^{1-\epsilon_4}} \right) \right]_{cd}^{ab} , \quad (\text{C.6})$$

$$\left[\left(\frac{\Pi^{\epsilon_1}}{\Pi^{\epsilon_2}} \right) \mathcal{M} \left(\frac{\Pi^{\epsilon_3}}{\Pi^{\epsilon_4}} \right) \right]_{cd}^{ab} = \left[\left(\frac{\Pi^{\epsilon_3}}{\Pi^{\epsilon_4}} \right) \mathcal{M} \left(\frac{\Pi^{\epsilon_1}}{\Pi^{\epsilon_2}} \right) \right]_{dc}^{ba} , \quad (\text{C.7})$$

$$\left[\left(\frac{\Pi^{\epsilon_1}}{\Pi^{\epsilon_2}} \right) \mathcal{M} \left(\frac{\Pi^{\epsilon_3}}{\Pi^{\epsilon_4}} \right) \right]_{cd}^{ab} \left[\left(\frac{\Pi^{\epsilon_2}}{\Pi^{\epsilon_1}} \right) \mathcal{M} \left(\frac{\Pi^{\epsilon_4}}{\Pi^{\epsilon_3}} \right) \right]_{ef}^{cd} = \delta_{ae} \delta_{bf} , \quad (\text{C.8})$$

where $\epsilon_i = 0, 1$ which correspond to not having and having a factor of Π respectively.

D On the resolution of the zero-mode problem

In section 4.1 the R-sector zero-mode spectrum in eqs.(4.31) has been obtained by requiring consistency with the relevant worldvolume interpretation. But it should also be possible to get the same result from the open string theory. As mentioned earlier it is not clear how to arrive at the result in eqs.(4.31) starting from the algebraic structure as given in eqs.(4.25, 4.28) and (4.29). Here we shall discuss certain curious features of the problem that might be relevant for the actual resolution. Notice that the spectrum in

eqs.(4.31) is exactly same as that in flat space where the definition of the ground states is known. If we follow the same definition then we have¹⁴:

$$S_0^a |I\rangle = \frac{1}{\sqrt{2}} \sigma_{a\dot{a}}^I |\dot{a}\rangle , \quad S_0^a |\dot{a}\rangle = \frac{1}{\sqrt{2}} \sigma_{a\dot{a}}^I |I\rangle . \quad (\text{D.1})$$

Then the problem will be to compute:

$$z_{a\dot{a}}^I := \langle I | \tilde{S}_0^a | \dot{a} \rangle . \quad (\text{D.2})$$

This can be done very easily: using eq.(4.30) and $\langle \dot{a} | H_0 | \dot{b} \rangle = 0$ one gets,

$$\sigma_{a\dot{a}}^I z_{a\dot{b}}^I = 0 . \quad (\text{D.3})$$

Then expanding $z_{a\dot{a}}^I$ in complete basis of odd rank $SO(8)$ gamma matrices:

$$z_{a\dot{a}}^I = c \sigma_{a\dot{a}}^I + c_{JK} \sigma_{a\dot{a}}^{IJK} , \quad (\text{D.4})$$

and substituting into eq.(D.3) one gets: $c = 0$, $c_{IJ} = 0$ implying,

$$z_{a\dot{a}}^I = 0 . \quad (\text{D.5})$$

There is also a way to get the same result using the existing algebraic structure without having any prior knowledge of H_0 eigenvalues. Using the first set of anti-commutators in eqs.(4.28) one obtains,

$$(\tilde{S}_0 \tilde{S}_0) = (S_0 S_0) = 4 , \quad (\tilde{S}_0^{IJKL} \tilde{S}) = (S \sigma^{IJKL} S) = 0 , \quad \forall I, J, K, L . \quad (\text{D.6})$$

Using these equations and the first relation in (4.25) one can show,

$$\tilde{S}_0^a = \frac{1}{56} \sum_{IJ} \hat{\lambda}_{IJ} (S \sigma^{IJ} S) (\sigma^{IJ} \tilde{S})^a . \quad (\text{D.7})$$

In this particular discussion we shall write the summations explicitly instead of following the repeated-index convention for them. At this stage we introduce the ground states as in eqs.(D.1). Then using eq.(D.7) one obtains:

$$z_{a\dot{a}}^K = \frac{1}{56} \sum_{IJLb} \hat{\lambda}_{IJ} \langle K | (S \sigma^{IJ} S) | L \rangle \sigma_{ab}^{IJ} z_{b\dot{a}}^L . \quad (\text{D.8})$$

¹⁴Without any loss of generality we have chosen $\beta = i/2\pi$ to avoid clutter in various expressions.

Using the result: $\langle K | (S\sigma^{IJ}S) | L \rangle = 4(\delta_{IK}\delta_{JL} - \delta_{IL}\delta_{JK})$ one finds,

$$\sum_{Jb} \left(7\delta^{IJ}\delta_{ab} - \hat{\lambda}_{IJ}\sigma_{ab}^{IJ} \right) z_{ba}^J = 0 . \quad (\text{D.9})$$

Then expanding $z_{a\dot{a}}^I$ as in eq.(D.4) and substituting this back into eq.(D.9) one gets an equation whose right hand side is zero and left hand side is expanded in terms of first and third rank $SO(8)$ gamma matrices. We then set the coefficient of each independent term to zero. For example, the coefficient of σ^I and σ^J ($J \neq I$) are given by $c\{8 - (8 - 2d)\hat{\lambda}_I\}$ and $-2c_{IJ}\{(8 - 2d)\hat{\lambda}_I - 1 - \hat{\lambda}_{IJ}\}$ respectively, where $d + 2 = p + 1$ is the dimension of worldvolume. This gives the same result as in eq.(D.2).

Given this result one may use the completeness relation: $\langle I | I \rangle + \langle \dot{a} | \dot{a} \rangle = 1$ to conclude that $\tilde{S}_0^a$ does not appear as a non-trivial operator in the theory. Notice that for the even D-branes and also for the non-BPS D-branes in flat space¹⁵ $\tilde{S}_0^a$ is completely eliminated classically so that it does not appear at all in the quantum theory. The above result seems to suggest that in our case $\tilde{S}_0^a$ is eliminated in the full quantum theory.

We have argued before that Δ^{ab} , as defined in (4.28), can not be a non-zero c-number, neither can it be fixed in the present theory. Notice that the above result explains this situation as it implies that an arbitrary matrix element of Δ^{ab} , which is potentially an operator, is always zero. However it is not clear to us how to interpret, for example, the non-zero anti-commutator of $\tilde{S}_0^a$ from the present point of view.

An argument using bosonisation

Here we shall attempt to understand the same result i.e. why $\tilde{S}_0^a$ may not appear as a non-trivial operator in the theory once the ground states are defined as in eqs.(D.1) in a different way using bosonisation. This approach is also not completely understood to us, but the following arguments show that it may be worth understanding this better.

Notice that the zero mode structure that we are dealing with, namely the first equation in (4.25), also appears in the discussion of non-BPS D-branes in flat background. Therefore it should be possible to understand these rules in that context itself. We therefore

¹⁵Open string spectrum for a non-BPS D-brane was obtained in [21] from the bi-local boundary condition without encountering the present problem. This is simply because in flat background the fermions are massless and there is no mixing between S and $\tilde{S}$ variables. Therefore all the $\tilde{S}$ variables, including the zero modes, are classically eliminated against the S variables. Because of this the operator $\tilde{S}_0^a$ does not appear at all in the open string theory.

consider the massless fermions for simplicity for the present argument. From the mode expansion in the bulk one gets,

$$S_0^a = \oint \frac{dz}{2\pi i} z^{-1/2} S^a(z) , \quad \tilde{S}_0^a = - \oint \frac{d\bar{z}}{2\pi i} \bar{z}^{-1/2} \tilde{S}^a(\bar{z}) , \quad (\text{D.10})$$

where the coordinates $(z, \bar{z})$ on the upper half plane (UHP) are related to the coordinates (τ, σ) on the strip (euclidean) as: $z = e^{\tau+i\sigma}$. For the operators in the bulk the contours in eqs.(D.10) do not pass through the real line. To define the above zero modes on the boundary one can go to the doubled surface by using the relevant boundary condition. This boundary condition needs to be linear as the right hand sides of eqs.(D.10) are linear in GS fermions. We already know that the boundary condition is actually quadratic. To see what kind of linear boundary conditions one might expect one can first bosonise $S^a(\tau, \sigma)$ and $\tilde{S}^a(\tau, \sigma)$ in terms of four left and right moving bosons: $\phi^i(\tau, \sigma)$ and $\tilde{\phi}^i(\tau, \sigma)$ respectively with $i = 1, \dots, 4$ [34, 35]. Then using the relevant boundary conditions relating ϕ and $\tilde{\phi}$ ¹⁶ and properly refermionising them back one will arrive at the desired expression. This is a detailed task by itself and we shall not attempt to do it explicitly here. But it seems plausible that the manifest $SO(8)$ covariance will require the final expression to be of the following form:

$$\tilde{S}^a(\tau, \sigma) = \widehat{\mathcal{M}}_{a\dot{a}} S^{\dot{a}}(\tau, \sigma) , \quad S^a(\tau, \sigma) = \widehat{\mathcal{M}}_{a\dot{a}} \tilde{S}^{\dot{a}}(\tau, \sigma) , \quad \text{at } \sigma = 0, \pi , \quad (\text{D.11})$$

where $\widehat{\mathcal{M}} = \sigma^{I_1 \dots I_{(p-1)}}$ is the spinor representation corresponding to the set of reflections given by $\hat{\lambda}_I$ in (4.27). Following [9, 36] one would then conclude that $S^{\dot{a}}(\tau, \sigma)$ and $\tilde{S}^{\dot{a}}(\tau, \sigma)$ are the spin fields that can be obtained by suitably bosonising and refermionising $S^a(\tau, \sigma)$ and $\tilde{S}^a(\tau, \sigma)$ respectively.

We shall now discuss the doubling trick in the present context. By using this trick one usually reduces an arbitrary correlator on the UHP to a holomorphic correlator on the full plane. This trick was suitably developed in [21] to handle the bi-local boundary conditions relevant to our case. It was shown that a correlator with an arbitrary number of bulk insertions on the UHP, but without any boundary insertions, can be mapped to a holomorphic correlator involving the holomorphic triad: $\{\mathcal{S}^a(z), \mathcal{S}^{\dot{a}}(z), \Psi^I(z)\}$. If the doubling trick works then the same should be true even in presence of the boundary

¹⁶Non-BPS D-branes were previously studied in [19] using such non-abelian bosonisation from the closed string point of view. Similar analysis can possibly be done in the open string picture to find boundary conditions for the bosons.

insertions. A suitable doubling that properly implements this is the following,

$$\mathcal{S}^a(\tau, \sigma) = \begin{cases} S^a(\tau, \sigma) & 0 \leq \sigma \leq \pi , \\ \widehat{\mathcal{M}}_{a\dot{a}} \tilde{S}^{\dot{a}}(\tau, 2\pi - \sigma) & \pi \leq \sigma \leq 2\pi . \end{cases} \quad (\text{D.12})$$

$$\mathcal{S}^{\dot{a}}(\tau, \sigma) = \begin{cases} S^{\dot{a}}(\tau, \sigma) & 0 \leq \sigma \leq \pi , \\ \widehat{\mathcal{M}}_{\dot{a}a}^{-1} \tilde{S}^a(\tau, 2\pi - \sigma) & \pi \leq \sigma \leq 2\pi . \end{cases} \quad (\text{D.13})$$

We can now express the zero modes on the boundary in the following way:

$$S_0^a = \oint \frac{dz}{2\pi i} z^{-1/2} \mathcal{S}^a(z) , \quad \tilde{S}_0^a = \widehat{\mathcal{M}}_{a\dot{a}} \oint \frac{dz}{2\pi i} z^{-1/2} \mathcal{S}^{\dot{a}}(z) , \quad (\text{D.14})$$

where the contours are understood to be around points on the real line.

Given eq.(D.14) we can now proceed to compute the matrix element in (D.2):

$$z_{a\dot{a}}^I = \lim_{w \rightarrow 0} \frac{1}{w} \widehat{\mathcal{M}}_{ab} \oint \frac{dz}{2\pi i} z^{-1/2} \langle \Psi^I(-1/w) \mathcal{S}^b(z) \mathcal{S}^{\dot{a}}(0) \rangle = 0 , \quad (\text{D.15})$$

as the holomorphic correlator itself is zero because of the momentum conservation of the bosons. Another way to understand why $\tilde{S}_0^a$ does not appear as a non-trivial operator is the following: Notice that the operators in (D.14) involve branch cuts. For any sensible computation these branch cuts need to be cancelled by other branch cuts originating from operators in the neighbourhood. It turns out that $\tilde{S}_0^a$ is not generically well defined in this sense in the neighbourhood of the ground states as defined in eqs.(D.1). For example, let us consider the following matrix element: $\langle \dot{b} | \tilde{S}_0^b \tilde{S}_0^a | \dot{a} \rangle$. Although the momentum conservation can be satisfied in this case, branch cuts are not removed:

$$\langle \dot{b} | \tilde{S}_0^b \tilde{S}_0^a | \dot{a} \rangle = \lim_{u \rightarrow 0} \frac{1}{u} \widehat{\mathcal{M}}_{b\dot{c}} \widehat{\mathcal{M}}_{a\dot{d}} \oint \frac{dz}{2\pi i} z^{-1/2} \oint \frac{dw}{2\pi i} w^{-1/2} \langle \mathcal{S}^b(-1/u) \mathcal{S}^{\dot{c}}(z) \mathcal{S}^{\dot{d}}(w) \mathcal{S}^{\dot{a}}(0) \rangle \quad (\text{D.16})$$

Contracting the w -contour to zero and using the OPE: $\mathcal{S}^{\dot{d}}(w) \mathcal{S}^{\dot{a}}(0) \sim \delta_{\dot{d}\dot{a}}/w$ one finds that the branch cut in w is not removed.

Although the above explanation may seem to be convincing, the present approach has to be understood better because of a potential caveat that we shall discuss now. If we have a linear boundary condition like in (D.11) then it is expected that the following bi-local boundary condition [21] will be reproduced:

$$\tilde{S}^a(\tau, \sigma) \tilde{S}^b(\tau', \sigma) = \widehat{\mathcal{M}}_{cd}^{ab} S^c(\tau, \sigma) S^d(\tau', \sigma) , \quad \text{at } \sigma = 0, \pi , \quad (\text{D.17})$$

where $\widehat{\mathcal{M}}_{cd}^{ab}$ is defined in eq.(4.26). This will be satisfied if the following equation holds as an equality of OPE's:

$$\widehat{\mathcal{M}}_{a\dot{a}}\widehat{\mathcal{M}}_{b\dot{b}}\mathcal{S}^{\dot{a}}(z)\mathcal{S}^{\dot{b}}(w) = \widehat{\mathcal{M}}_{cd}^{ab}\mathcal{S}^c(z)\mathcal{S}^d(w). \quad (\text{D.18})$$

This is almost satisfied, but not quite because of the subtlety involving fourth rank tensors. The fourth rank tensor constructed out of $\mathcal{S}^a(z)$ is always self-dual whereas that constructed out of $\mathcal{S}^{\dot{a}}(z)$ is always anti-self-dual. To see this more explicitly notice that the relevant OPEs can be computed using bosonisation as in, for example, [35]. The final results can be expanded in terms of the complete basis constructed out of $SO(8)$ gamma matrices. For the conjugate spinors one gets [35],

$$\begin{aligned} \mathcal{S}^{\dot{a}}(z)\mathcal{S}^{\dot{b}}(w) &= \delta_{\dot{a}\dot{b}} \sum_{n \geq -1} c_n^{(0)}(z-w)^n \mathcal{A}_n^{(0)}(\Psi(w)) + \bar{\sigma}_{\dot{a}\dot{b}}^{IJ} \sum_{n \geq 0} c_n^{(2)}(z-w)^n \mathcal{A}_n^{(2)IJ}(\Psi(w)) \\ &\quad + \bar{\sigma}_{\dot{a}\dot{b}}^{IJKL} \sum_{n \geq 1} c_n^{(4)}(z-w)^n \mathcal{A}_n^{(4)IJKL}(\Psi(w)), \\ &= \delta_{\dot{a}\dot{b}} \sum_{n \geq -1} c_n^{(0)}(z-w)^n \mathcal{A}_n^{(0)}(\Psi(w)) + \bar{\sigma}_{\dot{a}\dot{b}}^{IJ} \sum_{n \geq 0} c_n^{(2)}(z-w)^n \mathcal{A}_n^{(2)IJ}(\Psi(w)) \\ &\quad + \sum_{I,J,K,L \in \mathcal{K}^{(4)}} \bar{\sigma}_{\dot{a}\dot{b}}^{IJKL} \sum_{n \geq 1} c_n^{(4)}(z-w)^n \mathcal{A}_n^{(4,-)IJKL}(\Psi(w)), \end{aligned} \quad (\text{D.19})$$

where $\mathcal{A}_n^{(0)}$, $\mathcal{A}_n^{(2)}$ and $\mathcal{A}_n^{(4)}$ are anti-symmetric tensors of rank as indicated by the superscripts constructed using $\Psi^I(w)$ and its derivatives. Notice that in the first step the summation over the vector indices on the fourth rank tensors is free. But all $\bar{\sigma}^{IJKL}$ matrices are not independent. In the second step we have restricted this summation over only the independent ones¹⁷ using the anti-self-duality of the $\bar{\sigma}^{IJKL}$ matrices. $\mathcal{A}_n^{(4,-)IJKL} = \mathcal{A}_n^{(4)IJKL} - \frac{1}{4!}\epsilon^{IJKLMNPQ}\mathcal{A}_n^{(4)MNPQ}$ is therefore the anti-self-dual component of $\mathcal{A}_n^{(4)IJKL}(\Psi(w))$. Then using the following relations:

$$\left(\widehat{\mathcal{M}}\widehat{\mathcal{M}}^T\right)_{ab} = \delta_{ab}, \quad \left(\widehat{\mathcal{M}}\bar{\sigma}^{IJ}\widehat{\mathcal{M}}^T\right)_{ab} = \hat{\lambda}_{IJ}\sigma_{ab}^{IJ}, \quad \left(\widehat{\mathcal{M}}\bar{\sigma}^{IJKL}\widehat{\mathcal{M}}^T\right)_{ab} = \hat{\lambda}_{IJKL}\sigma_{ab}^{IJKL} \quad (\text{D.20})$$

one finds for the left hand side of eq(D.18),

$$\begin{aligned} \widehat{\mathcal{M}}_{a\dot{a}}\widehat{\mathcal{M}}_{b\dot{b}}\mathcal{S}^{\dot{a}}(z)\mathcal{S}^{\dot{b}}(w) &= \delta_{ab} \sum_{n \geq -1} c_n^{(0)}(z-w)^n \mathcal{A}_n^{(0)}(\Psi(w)) \\ &\quad + \hat{\lambda}_{IJ}\sigma_{ab}^{IJ} \sum_{n \geq 0} c_n^{(2)}(z-w)^n \mathcal{A}_n^{(2)IJ}(\Psi(w)) \end{aligned}$$

¹⁷This requires us to choose which ones we want to keep as independent. This is reflected in the choice of the set $\mathcal{K}^{(4)}$ that also appears in the bi-local boundary condition in (B.28, B.29).

$$+ \sum_{I,J,K,L \in \mathcal{K}_{(4)}} \hat{\lambda}_{IJKL} \sigma_{ab}^{IJKL} \sum_{n \geq 1} c_n^{(4)} (z-w)^n \mathcal{A}_n^{(4,-)IJKL}(\Psi(w)) , \quad (\text{D.21})$$

The OPE for the spinors is very similar to that in eq.(D.19) [35]:

$$\begin{aligned} \mathcal{S}^a(z) \mathcal{S}^b(w) &= \delta_{ab} \sum_{n \geq -1} c_n^{(0)} (z-w)^n \mathcal{A}_n^{(0)}(\Psi(w)) + \sigma_{ab}^{IJ} \sum_{n \geq 0} c_n^{(2)} (z-w)^n \mathcal{A}_n^{(2)IJ}(\Psi(w)) \\ &+ \sum_{I,J,K,L \in \mathcal{K}_{(4)}} \sigma_{ab}^{IJKL} \sum_{n \geq 1} c_n^{(4)} (z-w)^n \mathcal{A}_n^{(4,+)IJKL}(\Psi(w)) , \end{aligned} \quad (\text{D.22})$$

where in the last term the self-dual component: $\mathcal{A}_n^{(4,+)IJKL} = \mathcal{A}_n^{(4)IJKL} + \frac{1}{4!} \epsilon^{IJKLMNPQ} \mathcal{A}_n^{(4)MNPQ}$ appears. Then using various traces of the gamma matrices one shows:

$$\begin{aligned} \sum_{n \geq -1} c_n^{(0)} (z-w)^n \mathcal{A}_n^{(0)}(\Psi(w)) &= \frac{1}{8} \mathcal{S}^a(z) \mathcal{S}^a(w) , \\ \sum_{n \geq 0} c_n^{(2)} (z-w)^n \mathcal{A}_n^{(2)IJ}(\Psi(w)) &= \frac{1}{16} \sigma_{ab}^{IJ} \mathcal{S}^a(z) \mathcal{S}^b(w) , \\ \sum_{n \geq 1} c_n^{(4)} (z-w)^n \mathcal{A}_n^{(4,+)IJKL}(\Psi(w)) &= \frac{1}{192} \sigma_{ab}^{IJKL} \mathcal{S}^a(z) \mathcal{S}^b(w) . \end{aligned} \quad (\text{D.23})$$

Substituting these expressions back into eq.(D.21) one almost establishes eq.(D.18) except for the fact that the last equation in (D.23) tells us how to write self-dual tensors in terms of the spinors and not the ones which are anti-self-dual. It would be interesting to understand this problem better through an explicit bosonisation.

E A standard regularisation

Here we shall discuss a standard way to regularise the expression in (4.40). One may start with the following result derived in [37]:

$$\begin{aligned} &\pi^{-s/2} \Gamma(s/2) \sum_{n=-\infty}^{\infty} \left[\left(\frac{m}{\pi} \right)^2 + \left(\frac{n}{a} \right)^2 \right]^{-s/2} \\ &= \frac{am^{1-s}}{\pi^{(1-s)/2}} \left[\Gamma(s/2 - 1/2) + 2 \sum_{n=-\infty}^{\infty'} \frac{K_{1/2-s/2}(2m|an|)}{(m|an|)^{1/2-s/2}} \right] , \end{aligned} \quad (\text{E.1})$$

which is valid for negative s . The prime on the summation in the second term on the right hand side indicates that the $n = 0$ contribution is excluded. Setting $s = -1$ one can derive:

$$\sum_{n=1}^{\infty} \left[\left(\frac{n}{a} \right)^2 + \left(\frac{m}{2\pi} \right)^2 \right]^{1/2} = -\frac{am^2}{16\pi^2} \Gamma(-1) - \frac{m}{4\pi} - \frac{m}{2\pi^2} \sum_{n=1}^{\infty} \frac{1}{n} K_1(amn) , \quad (\text{E.2})$$

where $K_\nu(x)$ is a modified Bessel function. The Casimir energy of a one dimensional massive scalar field with mass m can be obtained from the above result where a is proportional to the size of the system. The first term on the right hand side is divergent and corresponds to a constant energy density. Dropping it off is the standard regularisation. The second term is independent of the size of the system - ignoring this is a constant shift of the total energy [37]. Therefore we may write:

$$\sum_{n=1}^{\infty} \left[\left(\frac{n}{a} \right)^2 + \left(\frac{m}{2\pi} \right)^2 \right]_{reg.}^{1/2} = -\frac{m}{2\pi^2} \sum_{n=1}^{\infty} \frac{1}{n} K_1(amn) , \quad (\text{E.3})$$

Now noticing that the right hand side of eq.(4.40) can be written in the following form,

$$h_0(m) = 8 \sum_{n=1}^{\infty} \sqrt{n^2 + \left(\frac{m}{2\pi} \right)^2} - 4 \sum_{n=1}^{\infty} \sqrt{\left(\frac{n}{2} \right)^2 + \left(\frac{m}{2\pi} \right)^2} , \quad (\text{E.4})$$

and using eq.(E.3) one gets the following result:

$$[h_0(m)]_{reg.} = \frac{2m}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n} [K_1(2mn) - 2K_1(mn)] . \quad (\text{E.5})$$

Notice that using the limiting behaviour: $\lim_{x \rightarrow 0} x K_1(x) \rightarrow 1$ one gets the correct massless limit:

$$\lim_{m \rightarrow 0} [h_0(m)]_{reg.} \rightarrow -\frac{1}{2} . \quad (\text{E.6})$$

References

- [1] A. Sen, “Tachyon dynamics in open string theory,” Int. J. Mod. Phys. A **20**, 5513 (2005) [arXiv:hep-th/0410103].
- [2] N. Drukker, D. J. Gross and N. Itzhaki, “Sphalerons, merons and unstable branes in AdS,” Phys. Rev. D **62**, 086007 (2000) [arXiv:hep-th/0004131]; K. Peeters and M. Zamaklar, “AdS/CFT description of D-particle decay,” Phys. Rev. D **71**, 026007 (2005) [arXiv:hep-th/0405125]; Y. L. He and P. Zhang, “ZZ brane decay in D dimensions,” JHEP **0611**, 014 (2006) [arXiv:hep-th/0607188]; D. Israel and E. Rabinovici, “Rolling tachyon in anti-de Sitter space-time,” arXiv:hep-th/0609087.
- [3] A. Sen, “Universality of the tachyon potential,” JHEP **9912**, 027 (1999) [arXiv:hep-th/9911116].

- [4] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2**, 231 (1998) [*Int. J. Theor. Phys.* **38**, 1113 (1999)] [arXiv:hep-th/9711200]; S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett. B* **428**, 105 (1998) [arXiv:hep-th/9802109]; E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2**, 253 (1998) [arXiv:hep-th/9802150].
- [5] M. Blau, J. Figueroa-O’Farrill, C. Hull and G. Papadopoulos, “A new maximally supersymmetric background of IIB superstring theory,” *JHEP* **0201**, 047 (2002) [arXiv:hep-th/0110242].
- [6] R. R. Metsaev, “Type IIB Green-Schwarz superstring in plane wave Ramond-Ramond background,” *Nucl. Phys. B* **625**, 70 (2002) [arXiv:hep-th/0112044].
- [7] R. R. Metsaev and A. A. Tseytlin, “Exactly solvable model of superstring in plane wave Ramond-Ramond background,” *Phys. Rev. D* **65**, 126004 (2002) [arXiv:hep-th/0202109].
- [8] D. Berenstein, J. M. Maldacena and H. Nastase, “Strings in flat space and pp waves from $N = 4$ super Yang Mills,” *JHEP* **0204**, 013 (2002) [arXiv:hep-th/0202021].
- [9] M. B. Green, J. H. Schwarz and E. Witten, “SUPERSTRING THEORY. VOL. 1: INTRODUCTION,”
- [10] J. Dai, R. G. Leigh and J. Polchinski, “NEW CONNECTIONS BETWEEN STRING THEORIES,” *Mod. Phys. Lett. A* **4**, 2073 (1989); J. Polchinski, S. Chaudhuri and C. V. Johnson, “Notes on D-Branes,” arXiv:hep-th/9602052; J. Polchinski, “Lectures on D-branes,” arXiv:hep-th/9611050;
- [11] A. Sen, “Non-BPS states and branes in string theory,” arXiv:hep-th/9904207; A. Lerda and R. Russo, “Stable non-BPS states in string theory: A pedagogical review,” *Int. J. Mod. Phys. A* **15**, 771 (2000) [arXiv:hep-th/9905006]; O. Bergman and M. R. Gaberdiel, “Non-BPS Dirichlet branes,” *Class. Quant. Grav.* **17**, 961 (2000) [arXiv:hep-th/9908126]; J. H. Schwarz, “TASI lectures on non-BPS D-brane systems,” arXiv:hep-th/9908144; M. R. Gaberdiel, “Lectures on non-BPS Dirichlet branes,” *Class. Quant. Grav.* **17**, 3483 (2000) [arXiv:hep-th/0005029].

- [12] A. Dabholkar and S. Parvizi, “Dp branes in pp-wave background,” Nucl. Phys. B **641**, 223 (2002) [arXiv:hep-th/0203231].
- [13] O. Bergman, M. R. Gaberdiel and M. B. Green, “D-brane interactions in type IIB plane-wave background,” JHEP **0303**, 002 (2003) [arXiv:hep-th/0205183]; M. R. Gaberdiel and M. B. Green, “The D-instanton and other supersymmetric D-branes in IIB plane-wave string Annals Phys. **307**, 147 (2003) [arXiv:hep-th/0211122]; M. R. Gaberdiel, M. B. Green, S. Schafer-Nameki and A. Sinha, “Oblique and curved D-branes in IIB plane-wave string theory,” JHEP **0310**, 052 (2003) [arXiv:hep-th/0306056].
- [14] K. Skenderis and M. Taylor, “Open strings in the plane wave background. I: Quantization and symmetries,” Nucl. Phys. B **665**, 3 (2003) [arXiv:hep-th/0211011]; K. Skenderis and M. Taylor, “Open strings in the plane wave background. II: Superalgebras and spectra,” JHEP **0307**, 006 (2003) [arXiv:hep-th/0212184]; K. Skenderis and M. Taylor, “Properties of branes in curved spacetimes,” JHEP **0402**, 030 (2004) [arXiv:hep-th/0311079].
- [15] M. Billo and I. Pesando, “Boundary states for GS superstrings in an Hpp wave background,” Phys. Lett. B **536**, 121 (2002) [arXiv:hep-th/0203028]; P. Bain, K. Peeters and M. Zamaklar, “D-branes in a plane wave from covariant open strings,” Phys. Rev. D **67**, 066001 (2003) [arXiv:hep-th/0208038]; Y. Hikida and S. Yamaguchi, “D-branes in pp-waves and massive theories on worldsheet with boundary,” JHEP **0301**, 072 (2003) [arXiv:hep-th/0210262]; J. w. Kim, B. H. Lee and H. S. Yang, “Superstrings and D-branes in plane wave,” Phys. Rev. D **68**, 026004 (2003) [arXiv:hep-th/0302060]; K. S. Cha, B. H. Lee and H. S. Yang, “Intersecting D-branes in type IIB plane wave background,” Phys. Rev. D **68**, 106004 (2003) [arXiv:hep-th/0307146]; C. Y. Park, “Open string spectrum in pp-wave background,” J. Korean Phys. Soc. **44**, 235 (2004) [arXiv:hep-th/0308151]; K. S. Cha, B. H. Lee and H. S. Yang, “A complete classification of D-branes in type IIB plane wave background,” JHEP **0403**, 058 (2004) [arXiv:hep-th/0310177]; M. Sakaguchi and K. Yoshida, “D-branes of covariant AdS superstrings,” Nucl. Phys. B **684**, 100 (2004) [arXiv:hep-th/0310228]; T. Mattik, “Branes in the plane wave background with gauge field condensates,” JHEP **0506**, 041 (2005) [arXiv:hep-th/0501088]; B. H. Lee, J. w. Lee, C. Park and H. S. Yang, “More on supersymmetric D-branes in type IIB plane wave background,”

- JHEP **0601**, 015 (2006) [arXiv:hep-th/0506091]; T. Mattik, “Boundary fermions and the plane wave,” JHEP **0605**, 045 (2006) [arXiv:hep-th/0603153].
- [16] K. Skenderis and M. Taylor, “Branes in AdS and pp-wave spacetimes,” JHEP **0206**, 025 (2002) [arXiv:hep-th/0204054]; P. Bain, P. Meessen and M. Zamaklar, “Supergravity solutions for D-branes in Hpp-wave backgrounds,” Class. Quant. Grav. **20**, 913 (2003) [arXiv:hep-th/0205106]; S. S. Pal, “Solution to worldvolume action of D3 brane in pp-wave background,” Mod. Phys. Lett. A **17**, 1735 (2002) [arXiv:hep-th/0205303]; R. R. Metsaev, “Supersymmetric D3 brane and $N = 4$ SYM actions in plane wave backgrounds,” Nucl. Phys. B **655**, 3 (2003) [arXiv:hep-th/0211178].
- [17] M. B. Green and M. Gutperle, “Light-cone supersymmetry and D-branes,” Nucl. Phys. B **476**, 484 (1996) [arXiv:hep-th/9604091].
- [18] N. D. Lambert and P. C. West, “D-branes in the Green-Schwarz formalism,” Phys. Lett. B **459**, 515 (1999) [arXiv:hep-th/9905031].
- [19] P. Mukhopadhyay, “Unstable non-BPS D-branes of type-II string theories in light-cone Green-Schwarz formalism,” Nucl. Phys. B **600**, 285 (2001) [arXiv:hep-th/0011047].
- [20] N. V. Suryanarayana, August 2004 - unpublished.
- [21] P. Mukhopadhyay, “Non-Bps D-Branes In Light-Cone Green-Schwarz Formalism,” JHEP **0501**, 059 (2005) [arXiv:hep-th/0411103].
- [22] T. Yoneya, “Spontaneously broken space-time supersymmetry in open string theory without GSO projection,” Nucl. Phys. B **576**, 219 (2000) [arXiv:hep-th/9912255].
- [23] N. Berkovits, “Super-Poincare covariant quantization of the superstring,” JHEP **0004**, 018 (2000) [arXiv:hep-th/0001035].
- [24] P. Mukhopadhyay, “On D-Brane Boundary State Analysis In Pure-Spinor Formalism,” JHEP **0603**, 066 (2006) [arXiv:hep-th/0505157].
- [25] P. Mukhopadhyay, “DDF construction and D-brane boundary states in pure spinor formalism,” JHEP **0605**, 055 (2006) [arXiv:hep-th/0512161].

- [26] P. Mukhopadhyay, “Tachyon condensation and ”non-BPS” D-branes in a Ramond-Ramond plane wave background,” Talk presented at Great Lakes Strings Conference, March-April, 2006.
- [27] P. Mukhopadhyay, “A Universality in PP-Waves,” JHEP **0706**, 061 (2007) [arXiv:0704.0085 [hep-th]].
- [28] A. Sen, “Supersymmetric world-volume action for non-BPS D-branes,” JHEP **9910**, 008 (1999) [arXiv:hep-th/9909062]. M. R. Garousi, “Tachyon couplings on non-BPS D-branes and Dirac-Born-Infeld action,” Nucl. Phys. B **584**, 284 (2000) [arXiv:hep-th/0003122]. E. A. Bergshoeff, M. de Roo, T. C. de Wit, E. Eyras and S. Panda, “T-duality and actions for non-BPS D-branes,” JHEP **0005**, 009 (2000) [arXiv:hep-th/0003221]; J. Kluson, “Proposal for non-BPS D-brane action,” Phys. Rev. D **62**, 126003 (2000) [arXiv:hep-th/0004106].
- [29] E. Witten, “D-branes and K-theory,” JHEP **9812**, 019 (1998) [arXiv:hep-th/9810188]; A. Sen, “BPS D-branes on non-supersymmetric cycles,” JHEP **9812**, 021 (1998) [arXiv:hep-th/9812031]; P. Horava, “Type IIA D-branes, K-theory, and matrix theory,” Adv. Theor. Math. Phys. **2**, 1373 (1999) [arXiv:hep-th/9812135]; A. Sen, “Non-BPS states and branes in string theory,” arXiv:hep-th/9904207.
- [30] G. T. Horowitz and A. R. Steif, “STRINGS IN STRONG GRAVITATIONAL FIELDS,” Phys. Rev. D **42**, 1950 (1990). R. E. Rudd, “Light cone gauge quantization of 2-D sigma models,” Nucl. Phys. B **427**, 81 (1994) [arXiv:hep-th/9402106].
- [31] J. M. Maldacena and L. Maoz, “Strings on pp-waves and massive two dimensional field theories,” JHEP **0212**, 046 (2002) [arXiv:hep-th/0207284];
- [32] J. G. Russo and A. A. Tseytlin, “Constant magnetic field in closed string theory: An Exactly solvable model,” Nucl. Phys. B **448**, 293 (1995) [arXiv:hep-th/9411099]; J. G. Russo and A. A. Tseytlin, “On solvable models of type IIB superstring in NS-NS and R-R plane wave backgrounds,” JHEP **0204**, 021 (2002) [arXiv:hep-th/0202179]; J. G. Russo and A. A. Tseytlin, “A class of exact pp-wave string models with interacting light-cone gauge actions,” JHEP **0209**, 035 (2002) [arXiv:hep-th/0208114];

- [33] G. Papadopoulos, J. G. Russo and A. A. Tseytlin, “Solvable model of strings in a time-dependent plane-wave background,” *Class. Quant. Grav.* **20**, 969 (2003) [arXiv:hep-th/0211289].
- [34] D. Friedan, E. J. Martinec and S. H. Shenker, “Conformal Invariance, Supersymmetry And String Theory,” *Nucl. Phys. B* **271**, 93 (1986).
- [35] V. A. Kostelecky, O. Lechtenfeld, W. Lerche, S. Samuel and S. Watamura, “CONFORMAL TECHNIQUES, BOSONIZATION AND TREE LEVEL STRING AMPLITUDES,” *Nucl. Phys. B* **288**, 173 (1987).
- [36] E. Witten, “D = 10 Superstring Theory,”
- [37] J. Ambjorn and S. Wolfram, “Properties Of The Vacuum. 1. Mechanical And Thermodynamic,” *Annals Phys.* **147**, 1 (1983).